arXiv:1701.05661v3 [math.AP] 15 Feb 2017

Quasi-periodic two-scale homogenisation and effective spatial dispersion in high-contrast media

Shane Cooper

February 16, 2017

Abstract

The convergence of spectra via two-scale convergence for double-porosity models is well known. A crucial assumption in these works is that the stiff component of the body forms a connected set. We show that under a relaxation of this assumption the (periodic) two-scale limit of the operator is insufficient to capture the full asymptotic spectral properties of high-contrast periodic media. Asymptotically, waves of all periods (or quasi-momenta) are shown to persist and an appropriate extension of the notion of two-scale convergence is introduced. As a result, homogenised limit equations with none trivially quasimomenta dependence are found as resolvent limits of the original operator family, resulting in limiting spectral behaviour with a rich dependence on quasimomenta.

1 Introduction

The model problem to study of time-harmonic waves, with frequency ω in media with microstructure is

$$-\mathrm{div}\big(a(\tfrac{x}{\varepsilon})\boldsymbol{\nabla} u\big) = \omega^2 u \qquad \text{in } \Omega$$

where the wave u represents the information being propagated, such as pressure in acoustics, deformation in elasticity or electromagnetic fields in electromagnetism[1] The physical nature of the waves under investigation depends on the choice of the domain Ω, with propagation and localisation being studied when Ω is the whole space and vibration modes under investigation for bounded Ω subject to specific boundary conditions. The coefficients a are periodic and characterise the microstructured nature of the media:

$$a(y) = \begin{cases} a_1(y), & y \in Q_1, \\ a_0(y), & y \in Q_0, \end{cases}$$

where a_0, a_1 are (the square-root of) the wave speeds of the individual constitutive material components that make up the composite, see figure 1. The parameter ε represents the ratio between the observable length scale and the size of the microstructure and is typically small. Studying the asymptotic behaviour of these waves in the limit of vanishing ε, for fixed frequency, is important in applications and the homogenisation theorem, also called the long-wavelength or quasi-static approximation depending on the community, states that for fixed $\lambda = \omega^2$ the wave behaviour of a microstructured media for small enough ε can be approximated by studying the vibrations of an 'effective' homogeneous media whose wave speed a^{hom} is constant and determined directly from the 'local periodic' behaviour of the problem.

It has been known for some time now that the introduction of so-called high contrasts in the coefficients results in unusual effects in microstructured media such as memory effects and other non-local effects (e.g. [4, 6–8, 13, 14, 16, 26]). Those high-contrast problems of particular interest in the context of waves are the so-called double porosity models where the coefficients of the (inclusion) phase Q_0 are 'critically' scaled

[1] In elasticity and electromagnetism the wave equation describes certain polarised waves: e.g. Shear polarised wave in elasticity or Transverse Electric and Transverse Magnetic polarised waves for the Maxwell system.



Figure 1: A typical schematic of some three-dimensional composite media with period microstructure.

by taking them to be of the order ε^2. Physically, this critical scaling corresponds to the scale of the wave-length of u remaining 'long' within the matrix Q_1 but in Q_0 the wavelength is at the 'resonant' scale, i.e. of the same order as the size of the microstructure, thus violating the underlying intuition for the long-wave length approximation. V. Zhikov [28, 29] analysed the double porosity spectral problems in bounded [28] and unbounded [29] domains by developing the homogenisation theory for generalised double porosity-type problems within the framework of two-scale convergence of G. Nguetseng-G. Allaire [1, 25]. Zhikov derived two-scale limit spectral equations that contain a non-trivial coupling between micro- and macro-scales, and the de-coupling of these scales, which is possible due to certain underlying geometric assumptions, leads to an eigenvalue problem with a highly non-linear dependence described by a function $\beta(\lambda)$ of the spectral parameter $\lambda = \omega^2$. This β function describes in an explicit way the presence of strong band-gaps in applications to wave propagation: frequency bands where no modes can propagate within the medium, [17, 29], and has recently been shown to describe the asymptotic structure of the spectral band functions [12]. This Zhikov β function was later interpreted in the works of G. Bouchitté and D. Felbacq [5], in the context of TM polarised electromagnetic waves in a dilute dielectric two-dimensional photonic crystal fibre, as negative effective magnetism. In the context of elasticity, a matrix analogue of the β function is derived and plays the role of effective density that is frequency dependent [2, 3, 30]. Aside from the strong spectral band-gaps mentioned above this leads to "weak" gaps where reduced numbers of modes are allowed to propagate within the medium. The use of high-contrast media as wave-guides due to the presence of gaps in the spectrum of high-contrast media was studied using multi-scale asymptotics in [19] and supplemented with analysis based on two-scale convergence in [9].

The idea that high-contrast media can result in the appearance of non-standard constitutive laws and give rise to composite media with complex wave phenomena near micro-resonances has prompted a recent energetic pursuit of non-standard constitutive laws in the contexts of elasticity (e.g. [22]) and electromagnetism (e.g. [11, 21]) with applications in areas such as cloaking (e.g. [23, 24]). It was shown in the work of V. Smyshlyaev [27], building on related ideas in [13], that by increasing the elastic anisotropy of composite media and by considering "inter-connected" inclusions then the two-scale homogenised limit contain not only the temporal non-locality given by the Zhikov β function but also exhibit spatial non-locality. The presence of which leads to the phenomena of 'directional' localisation: the number of admissible propagating wave modes depends not only on the frequency but on the direction of propagation. These works have very recently been complemented with a systematic study containing rigorous asymptotic and spectral analysis of general mathematical constructions containing 'high-contrasts' [20]. Analysis based on the work [20] has led to the demonstration that the two-scale convergence is insufficient to fully study the spectrum of general high-contrast problems, see [15, Chapter 5], [10]. The reason for this inconsistency is due to the

presence of quasi-periodic microscopic oscillations that, while are asymptotically negligible due to geometric considerations in previously studied double porosity models, persist at leading-order in general high-contrast media.

The aim of this work is to two-fold. First we show that by a simple relaxing of the geometric assumptions in the double porosity model, namely the connectedness of the matrix or 'stiff' phase, leads to two-scale homogenised models that contain a new feature: the effective wave speed depending on the quasi-momenta in a highly discontinuous fashion, i.e. the non-standard constitutive equations for such high-contrast media exhibit spatial dispersion. The addition of this feature, on top of the temporal non-locality demonstrated by Zhikov, and the spatial non-locality demonstrated by Smyshlyaev, is due to the additional asymptotic presence of quasi-periodic waves on the microscale. The second aim of this work is to appropriately develop the two-scale frame-work, laid out in [20], for the study of quasi-periodic micro-oscillations in a generalised class of high-contrast problems. This framework will be used here to perform a spectral analysis of novel high-contrast models and additionally explain why such micro-oscillations were not present in the numerous previous works on high-contrast spectral and dynamic problems.

We end the introduction with some words on the **notation** used in this article. Vectors and vector-valued functions are represented by lower-case boldface symbols with the exception of the co-ordinate points. $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ denotes the Euclidean basis in $\mathbb{R}^3$. For a vector $\mathbf{u} \in \mathbb{R}^3$, we denote by u_i its component with respect to $\mathbf{e}_i$, and write

$$\mathbf{u} = (u_1, u_2, u_3) = \sum_{i=1}^{3} u_i \mathbf{e}_i.$$

Points in $\mathbb{R}^3$ will be denoted by the symbol x and points in the unit cell $Q := (0,1)^3$ will be denoted by y. The notation ∂_i will be used to denote partial differentiation with respect to the i-th coordinate variable, and we shall replace the suffix i with x_i or y_i when we wish to emphasis the macroscopic or microscopic variable. Similarly, the notion div_x, div_y, $\boldsymbol{\nabla}_x$, and $\boldsymbol{\nabla}_y$, are used for the divergence or gradient of a function in terms of x or y. The Einstein summation convention will not be used in this article, that is we do not sum with respect to repeated indices.

2 Problem formulation and Homogenisation

In this article we are concerned with the asymptotic analysis of the resolvent problem

$$\begin{cases} \text{Find } u_\varepsilon \in H_0^1(\Omega) \text{ such that} \\ -\operatorname{div}\big(a_\varepsilon(\frac{x}{\varepsilon})\boldsymbol{\nabla} u_\varepsilon\big) + u_\varepsilon = f_\varepsilon \qquad \text{in } \Omega \end{cases} \tag{2.1}$$

where $\varepsilon < 1$ is a small parameter, Ω is a smooth open bounded star-shaped domain[2] and $f_\varepsilon \in L^2(\Omega)$ known. The coefficient a_ε is given by

$$a_\varepsilon(y) = \begin{cases} a_1(y), & y \in Q_1, \\ \varepsilon^2 a_0(y), & y \in Q_0, \end{cases} \quad 0 \leq a_i, a_i^{-1} \in L^\infty(Q_i),\ a_i = 0 \text{ on } Q_{1-i}, i = 1, 2, \tag{2.2}$$

and the regions Q_0 and Q_1 are described as follows.

Geometric assumptions. For $j = 1, 2, 3$ we consider smooth domains S_j compactly contained in $(0,1)^2$ that have mutually disjoint closures. We denote by C_j the cylinder aligned to the j-th co-ordinate axis with cross-section S_j, i.e. $C_1 := \{y \in (0,1)^3 \,\big|\, y \in (0,1) \times S_1\}$, $C_2 := \{y \in (0,1)^3 \,\big|\, y = (z_2, z_3, z_1), z \in (0,1) \times S_2\}$ and $C_3 := \{y \in (0,1)^3 \,\big|\, y = (z_3, z_1, z_2), z \in (0,1) \times S_3\}$.

Then, for a given non-empty subset $\mathcal{I}$ of $\{1,2,3\}$, we consider $Q_1 = \cup_{i \in \mathcal{I}} C_i$. We denote by $\Gamma_i = \partial C_i \backslash \partial Q$ and $\Gamma = \bigcup_{i \in \mathcal{I}} \Gamma_i = \partial Q_1 \backslash \partial Q$.

[2] All the results and proofs follow through in an identical manner for the case where Ω is the whole space.



Figure 2: a) An example stiff component Q_1 consisting of one cylinder C_1 extending in the x_1 direction, i.e. $\mathcal{I} = \{1\}$. b) The stiff component Q_1 consists of two disjoint cylinders extending in the co-ordinate directions x_1, and x_3, i.e. $\mathcal{I} = \{1, 3\}$. c) Q_1 consists of mutually disjoint cylinders extending in all co-ordinate directions, i.e. $\mathcal{I} = \{1, 2, 3\}$.

Under this geometric assumption we determine for each $\boldsymbol{\theta} \in [0, 2\pi)^3$ the strong $\boldsymbol{\theta}$-quasi-periodic two-scale resolvent limit, cf. Section 5, of the self-adjoint operator A_ε associated to resolvent problem (2.1). That is, for a fixed $\boldsymbol{\theta} \in [0, 2\pi)^3$ and a given bounded sequence $f_\varepsilon \in L^2(\Omega)$ such that $f_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} f$, i.e.

$$\lim_{\varepsilon\to 0}\int_\Omega f_\varepsilon(x)\overline{\phi(x)\varphi\left(\tfrac{x}{\varepsilon}\right)}\,\mathrm{d}x = \int_\Omega\int_\square f(x,y)\overline{\phi(x)\varphi(y)}\,\mathrm{d}y\mathrm{d}x,$$
$$\forall \phi \in C_0^\infty(\Omega),\ \forall \varphi \in C^\infty_{\boldsymbol{\theta}}(\square)$$

we aim to determine the $\boldsymbol{\theta}$-quasi-periodic two-scale limit behaviour of the solution $u_\varepsilon \in H_0^1(\Omega)$ to

$$\int_\Omega \left(a_1(\tfrac{x}{\varepsilon}) + \varepsilon^2 a_0(\tfrac{x}{\varepsilon})\right)\boldsymbol{\nabla} u_\varepsilon(x)\cdot\overline{\boldsymbol{\nabla}\phi}(x)\,\mathrm{d}x \ + \int_\Omega u_\varepsilon(x)\overline{\phi}(x)\,\mathrm{d}x = \int_\Omega f_\varepsilon(x)\overline{\phi}(x)\,\mathrm{d}x, \qquad \forall \phi \in C_0^\infty(\Omega). \tag{2.3}$$

As f_ε is bounded in $L^2(\Omega)$, upon setting $\phi = u_\varepsilon$ in (2.3) we deduce that the sequences

$$||\sqrt{a_1}(\tfrac{\cdot}{\varepsilon})\boldsymbol{\nabla} u_\varepsilon||_{L^2(\Omega;\mathbb{C}^3)}, \quad ||\varepsilon\boldsymbol{\nabla} u_\varepsilon||_{L^2(\Omega;\mathbb{C}^3)}, \quad \text{and} \quad ||u_\varepsilon||_{L^2(\Omega)}, \tag{2.4}$$

are bounded. Let us describe the $\boldsymbol{\theta}$-quasi-periodic two-scale limit, referring to Section 6 for the details. The limit of $u_\varepsilon(x)$ will be a function $u(x, y)$, of two variables $x \in \Omega$, $y \in Q$, that is $\boldsymbol{\theta}$-quasi-periodic with respect to the second variable y. Furthermore, due to the fact that in each cylinder C_i, $i \in \mathcal{I}$, the gradient of u_ε is bounded, the limit u necessarily belongs to the (Bochner) space $L^2(\Omega; V_{\boldsymbol{\theta}})$ where

$$V_{\boldsymbol{\theta}} := \{v \in H^1_{\boldsymbol{\theta}}(Q) \,|\, \ v \text{ is constant in } C_i \text{ for each } i \in \mathcal{I}\}. \tag{2.5}$$

It follows from this that u is non-zero in cylinder C_i if and only if the i-th component θ_i of $\boldsymbol{\theta}$ is zero. If $\theta_i = 0$ then generally u_i is not only non-trivial but it is actually more regular in the x_i-th coordinate direction: $\partial_{x_i} u_i \in L^2(\Omega)$.
More precisely, for $\mathcal{I}_{\boldsymbol{\theta}} := \{i \in \mathcal{I} \,|\, \theta_i = 0\}$, we denote $L_{\boldsymbol{\theta}}$ as the closed subspace of $\mathbb{C}^3$ spanned by $\{\mathbf{e}_i\}_{i\in\mathcal{I}_{\boldsymbol{\theta}}}$, and show that the function u belongs to the set

$$\begin{aligned} U_{\boldsymbol{\theta}} = \big\{ u \in L^2(\Omega; H^1_{\boldsymbol{\theta}}(Q)) \,\big|\, & u = u_i \text{ on } \Omega\times C_i, \\ & \text{for some } \mathbf{u} \in L^2(\Omega; L_{\boldsymbol{\theta}}) \text{ with } \partial_i u_i \in L^2(\Omega) \text{ and } u_i\nu_i = 0 \text{ on } \partial\Omega \ \big\}, \end{aligned} \tag{2.6}$$

which is clearly a Hilbert space when endowed with the inner product

$$(u,v)_{U_{\boldsymbol{\theta}}} := \sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}}\int_\Omega \partial_i u_i(x)\overline{\partial_i v_i(x)}\,\mathrm{d}x + \int_\Omega\int_{Q_0} \boldsymbol{\nabla}_y u(x,y)\cdot\overline{\boldsymbol{\nabla}_y v(x,y)}\,\mathrm{d}y\mathrm{d}x$$
$$+ \int_\Omega\int_Q u(x,y)\overline{v(x,y)},\mathrm{d}y\mathrm{d}x.$$

Here, $\boldsymbol{\nu}$ is the outer unit normal to $\partial\Omega$.
For each fibre C_i there corresponds an effective constant material parameter $a_i^{\mathrm{hom}} > 0$ given by

$$a_i^{\mathrm{hom}} = \int_{C_i} a_1(y)[\partial_{y_i} N^{(i)}(y) + 1]\,\mathrm{d}y, \tag{2.7}$$

where $N^{(i)} \in H^1_{\#_i}(C_i) := \{u \in H^1(C_i) \,|\, u \text{ is 1-periodic in the variable } y_i\}$ is the unique solution to the cell problem

$$\begin{cases} \displaystyle\int_{C_i} a_1(y)\big[\boldsymbol{\nabla} N^{(i)}(y) + \mathbf{e}_i\big] \cdot \overline{\boldsymbol{\nabla}\phi}(y)\,\mathrm{d}y = 0, \qquad \forall \phi \in H^1_{\#_i}(C_i), \\ \displaystyle\int_{C_i} N^{(i)} = 0. \end{cases} \tag{2.8}$$

Then, for each $\boldsymbol{\theta} \in [0, 2\pi)^3$, the $\boldsymbol{\theta}$-quasi-periodic two-scale limiting problem is formulated as follows: For $f \in L^2(\Omega \times Q)$ find $u \in U_{\boldsymbol{\theta}}$ such that

$$\begin{aligned} \sum_{i \in \mathcal{I}_{\boldsymbol{\theta}}} \int_\Omega a_i^{\mathrm{hom}} \partial_{x_i} u_i(x) \overline{\partial_{x_i}\phi_i}(x)\,\mathrm{d}x + \int_\Omega \int_{Q_0} a_0(y) \boldsymbol{\nabla}_y u(x,y) \cdot \overline{\boldsymbol{\nabla}_y \phi(x,y)}\,\mathrm{d}y\mathrm{d}x \\ + \int_\Omega \int_Q u(x,y)\overline{\phi(x,y)}\,\mathrm{d}y\mathrm{d}x = \int_\Omega \int_Q f(x,y)\overline{\phi(x,y)}\,\mathrm{d}y\mathrm{d}x, \qquad \forall \phi \in U_{\boldsymbol{\theta}}. \end{aligned} \tag{2.9}$$

As a_i^{hom} are positive numbers and $a_0^{-1} \in L^\infty(Q_0)$ it follows that the left-hand side of the above problem defines an equivalent inner product on the space $U_{\boldsymbol{\theta}}$, and consequently the existence and uniqueness of solutions u to (2.9) are ensured by the Riesz representation theorem.
Setting $\phi = 0$ on Q_1 in (2.9) gives the equation

$$-\mathrm{div}_y\big(a_0(y)\boldsymbol{\nabla}_y u(x,y)\big) + u(x,y) = f(x,y), \qquad x \in \Omega,\ y \in Q_0,$$

and a subsequent integration by parts in (2.9) leads to the variational formula

$$\begin{aligned} \sum_{i \in \mathcal{I}_{\boldsymbol{\theta}}} \int_\Omega a_i^{\mathrm{hom}} \partial_{x_i} u_i(x) \overline{\partial_{x_i}\phi_i}(x)\,\mathrm{d}x + \sum_{i \in \mathcal{I}_{\boldsymbol{\theta}}} \int_\Omega \left(\int_{\Gamma_i} a_0(y) \boldsymbol{\nabla}_y u(x,y) \cdot \mathbf{n}(y)\,\mathrm{d}y \right) \overline{\phi_i(x)}\mathrm{d}x \\ + \sum_{i \in \mathcal{I}_{\boldsymbol{\theta}}} |C_i| \int_\Omega u_i(x)\overline{\phi_i(x)}\mathrm{d}x = \sum_{i \in \mathcal{I}_{\boldsymbol{\theta}}} \int_\Omega \left(\int_{C_i} f(x,y)\,\mathrm{d}y \right) \overline{\phi_i(x)}\mathrm{d}x, \end{aligned}$$

for all $\boldsymbol{\phi} \in L^2(\Omega, L_{\boldsymbol{\theta}})$, such that $\partial_i \phi_i \in L^2(\Omega)$. Sequentially setting, for each $j \in \mathcal{I}_{\boldsymbol{\theta}}$, $\phi_i = \delta_{ij}\phi$, $\phi \in C_0^\infty(\Omega)$, leads to the $\boldsymbol{\theta}$-quasi-periodic two-scale homogenised system of equations

$$\begin{cases} -a_i^{\mathrm{hom}} \partial_{x_i}^2 u_i(x) + \mathcal{T}_i(u) + |C_i| u_i = \langle f \rangle_i(x), & x \in \Omega, \\ -\mathrm{div}_y\big(a_0(y)\boldsymbol{\nabla}_y u(x,y)\big) + u(x,y) = f(x,y), & x \in \Omega,\ y \in Q_0, \\ u_0 = u_i \quad \text{on } \Omega \times \Gamma_i, \qquad u_i \nu_i = 0 \quad \text{on } \partial\Omega, & i \in \mathcal{I}. \end{cases} \tag{2.10}$$

Here

$$\mathcal{T}_i(u)(x) = \int_{\Gamma_i} a_0(y) \boldsymbol{\nabla}_y u(x,y) \cdot \mathbf{n}(y)\,\mathrm{d}S(y), \qquad \langle f \rangle_i(x) = \int_{C_i} f(x,y)\,\mathrm{d}y,$$

for $\mathbf{n}$ the outer unit normal of $\Gamma_i = \partial C_i \backslash \partial Q$. We now state the main result of the article.

Theorem 1. *Consider $f_\varepsilon \in L^2(\Omega)$, $f \in L^2(\Omega \times Q)$ such that $f_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} f$, and u_ε the solution to* (2.3). *Then u_ε converges, up to some subsequence, in the $\boldsymbol{\theta}$-quasi-periodic sense to $u \in U_{\boldsymbol{\theta}}$ the unique solution to* (2.9), *equivalently* (2.10).

An immediate consequence of Theorem 1 is that for each $\boldsymbol{\theta} \in [0,2\pi)^3$, the operator A_ε strong $\boldsymbol{\theta}$-quasi-periodically two-scale resolvent converges to the operator $A_{\boldsymbol{\theta}}^{\hom}$ associated to problem (2.10), see Section 5 for details. Consequently, Proposition 6, therein, informs us that the lower semi-continuity of the spectra in the Hausdorff sense is ensured:

$$\text{for every } \lambda \in \bigcup_{\boldsymbol{\theta}\in[0,2\pi)^3} \sigma(A_{\boldsymbol{\theta}}^{\hom}) \ \exists\, \lambda_\varepsilon \in \sigma(A_\varepsilon) \text{ such that } \lim_{\varepsilon\to 0} \lambda_\varepsilon = \lambda.$$

The structure of the limit spectrum $\bigcup_{\boldsymbol{\theta}\in[0,2\pi)^3} \sigma(A_{\boldsymbol{\theta}}^{\hom})$ is analysed in Section 4. In particular, we show that

Proposition 1. *The spectrum of $\bigcup_{\boldsymbol{\theta}\in[0,2\pi)^3} \sigma(A_{\boldsymbol{\theta}}^{\hom})$ is the union of the following two sets:*

- *The pure Bloch spectrum:*

$$\bigcup_{\boldsymbol{\theta}\in[0,2\pi)^3} \sigma(B_{\boldsymbol{\theta}}) = \sum_{m\in\mathbb{N}} \left[\min_{\boldsymbol{\theta}\in[0,2\pi)^3} \left(\mu_{\boldsymbol{\theta}}^{(m)}\right), \max_{\boldsymbol{\theta}\in[0,2\pi)^3} \left(\mu_{\boldsymbol{\theta}}^{(m)}\right) \right],$$

  where $\mu_{\boldsymbol{\theta}}^{(m)}$ are the eigenvalues of $B_{\boldsymbol{\theta}}$, see Section 3.2, ordered according to the min-max principle.

- *The spatial spectrum: $\{\lambda \in [0,\infty) \,|\, \gamma(\boldsymbol{\theta})(\lambda) \in \sigma(A_{\boldsymbol{\theta}})\}$, where $A_{\boldsymbol{\theta}}$ is introduced in Section 3.1 is an operator with compact resolvent. Here $\gamma : \mathbb{R}^3 \to \mathbb{S}^3$ is for each $\boldsymbol{\theta}$ a (possibly) sign-indefinite symmetric matrix defined by setting for $i \notin \mathcal{I}_{\boldsymbol{\theta}}$, $\gamma_{ij}(\boldsymbol{\theta}) = 0$ for all j, and $\gamma_{ij}(\boldsymbol{\theta}) = \beta_{\boldsymbol{\theta}}^{(ij)}$ otherwise, see* (4.6).

The above result claims that in general the limit spectrum is of band-gap structure with (possibly embedded and accumulating) point-spectrum. The spectral bands are prescribed by the spectrum of the pure Bloch operator $B_{\boldsymbol{\theta}}$ and the point spectrum is described by the non-linear eigenvalue problem $A(\boldsymbol{\theta})w = \gamma(\boldsymbol{\theta})(\lambda)w$.

Remark 1. In the double porosity case where Q_0 is a compactly contained subset of Q_1, it was demonstrated by V. Zhikov in [29] that the corresponding spectrum $\sigma(A_\varepsilon)$ Hausdorff converges to the limit spectrum of the periodic two-scale limit. From the context of the framework presented in this article, this result seems somewhat surprising due to the possible additional limit spectrum for $\boldsymbol{\theta} \in (0,2\pi)^3$. Yet, this is explained by noting that $V_{\boldsymbol{\theta}} = H_0^1(Q_0)$ for $\boldsymbol{\theta} \neq 0$ (recall (2.5)) and one finds $B_{\boldsymbol{\theta}}$ is independent of $\boldsymbol{\theta}$ for $\boldsymbol{\theta} \neq 0$. Furthermore, the pure Bloch spectrum is discrete here and consists precisely of the resonant frequencies of the inclusion Q_0.
Similarly, in the classical homogenisation regime when $Q_0 = \emptyset$, $Q_1 = Q$, then $V_{\boldsymbol{\theta}} = \{0\}$ for $\boldsymbol{\theta} \neq 0$. This explains why the role of periodic functions are of special importance in the classical and standard double-porosity contexts.

Remark 2. A seperate issue, not explored here, is the so-called spectral completeness statement, i.e. the question of whether or not the remaining criterion for Hausdorff convergence of spectra is satisfied: does it follow that

$$\text{for every } \lambda_\varepsilon \text{ such that } \lim_{\varepsilon\to 0} \lambda_\varepsilon = \lambda, \text{ then } \lambda \in \bigcup_{\boldsymbol{\theta}\in[0,2\pi)^3} \sigma(A_{\boldsymbol{\theta}}^{\hom}) \ ?$$

In general this will not be true due to the presence of the boundary, and the fact that Q_0 intersects the boundary, leads to the expectation that there exists non-trivial spectrum due to surface waves asymptotically localised near the boundary. For the case of Ω being the torus or the whole space the above assertion is expected to hold and will be explored in future works.

3 Quasi-periodic two-scale limit operator and resolvent convergence

For $\boldsymbol{\theta} \in [0,2\pi)^3$, we consider the subspace H which is the closure of $U_{\boldsymbol{\theta}}$ in $L^2(\Omega \times Q)$, i.e.

$$H = \big\{ u \in L^2(\Omega \times Q) \,\big|\, u = u_i \text{ on } \Omega \times C_i \quad \text{for some } \mathbf{u} \in L^2(\Omega; L_{\boldsymbol{\theta}}) \ \big\}.$$

Indeed, for $f \in H$, we have $f = f_i$ on $\Omega \times C_i$, $i \in \mathcal{I}$, and consequently we deduce that

$$||f||^2_{L^2(\Omega;L^2(Q_1))} = \sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}} |C_i|||f_i||^2_{L^2(\Omega)},$$

and therefore, H is closed in $L^2(\mathbb{R}^3;L^2(Q))$. It is also straight forward to show that $U_{\boldsymbol{\theta}}$ is dense in H. Defining on $U_{\boldsymbol{\theta}}$ the form

$$Q_{\boldsymbol{\theta}}(u,v) := \sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}} \int_\Omega a_i^{\hom} \partial_{x_i} u_i(x) \overline{\partial_{x_i} v_i(x)}\,\mathrm{d}x + \int_\Omega \int_{Q_0} a_0(y) \boldsymbol{\nabla}_y u(x,y) \cdot \overline{\boldsymbol{\nabla}_y v(x,y)}\,\mathrm{d}y\mathrm{d}x,$$

we find that, since $a_i^{\hom}$ are positive constants and $a_0^{-1} \in L^\infty(Q_0)$, $Q_{\boldsymbol{\theta}}$ is closed when considered as a form on H. Setting $A_{\boldsymbol{\theta}}^{\hom} : D(A_{\boldsymbol{\theta}}^{\hom}) \subset H \to H$ to be the unbounded self-adjoint operator generated by $Q_{\boldsymbol{\theta}}$, for $f \in L^2(\mathbb{R}^3;Q)$ the $\boldsymbol{\theta}$-quasi-periodic two-scale homogenised limit problem (2.9) takes the form $A_{\boldsymbol{\theta}}^{\hom} u = P_{\boldsymbol{\theta}} f$. Here, $P_{\boldsymbol{\theta}} : L^2(\mathbb{R}^3;L^2(Q)) \to H$ is the orthogonal projection given by

$$P_{\boldsymbol{\theta}} f(x,y) = \begin{cases} \int_{C_i} f(x,y)\,\mathrm{d}y, & y \in C_i, \\ f(x,y), & y \in Q_0. \end{cases}$$

An immediate consequence of Theorem 1 is that for each $\boldsymbol{\theta} \in [0,2\pi)^3$, the operator A_ε strong $\boldsymbol{\theta}$-quasi-periodic two-scale resolvent converges to $A_{\boldsymbol{\theta}}^{\hom}$, see Section 5, definition 2.

3.1 Spatial operators

Introducing the notation

$$D := \begin{pmatrix} \partial_{x_1} & 0 & 0 \\ 0 & \partial_{x_2} & 0 \\ 0 & 0 & \partial_{x_3} \end{pmatrix}, \qquad A^{\hom} := \begin{pmatrix} a_1^{\hom} & 0 & 0 \\ 0 & a_2^{\hom} & 0 \\ 0 & 0 & a_3^{\hom} \end{pmatrix},$$

we consider the Hilbert space

$$H_{\boldsymbol{\theta}} := \left\{ \mathbf{u} \in L^2(\Omega;L_{\boldsymbol{\theta}}) \,|\, D\mathbf{u} \in L^2(\Omega), u_i\nu_i = 0 \text{ on } \partial\Omega, i = 1,2,3 \right\},$$

endowed with the inner product

$$(u,v)_H := \int_\Omega D\mathbf{u} \cdot \overline{D\mathbf{v}},$$

and the following bilinear form defined on H:

$$\alpha_{\boldsymbol{\theta}}(u,v) := \sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}} \int_\Omega a_i^{\hom} \partial_{x_i} u_i \cdot \overline{\partial_{x_i} v_i} = \int_\Omega A^{\hom} D\mathbf{u} \cdot \overline{D\mathbf{v}}, \qquad \mathbf{u},\mathbf{v} \in H_{\boldsymbol{\theta}}.$$

Note that for $\boldsymbol{\theta}$ such that $\mathcal{I}_{\boldsymbol{\theta}} = \{i \in \mathcal{I} \,|\, \theta_i = 0\} = \emptyset$ then $H_{\boldsymbol{\theta}}$ is zero and for such $\boldsymbol{\theta}$ we define our 'spatial' operator $A_{\boldsymbol{\theta}}$ to be the zero map. Otherwise, $\alpha_{\boldsymbol{\theta}}$ is a positive form on $H_{\boldsymbol{\theta}}$ and therefore has a positive self-adjoint operator $A_{\boldsymbol{\theta}}$, densely defined in $L^2(\Omega;Ł_{\boldsymbol{\theta}})$, associated with the form. The space H is compactly embedded[3] into L^2, and consequently the spatial operator $A_{\boldsymbol{\theta}}$ has compact resolvent and therefore its spectrum is discrete.

[3]This follows from an application of Vitali's theorem, which is permissible by noting that since u_i has an L^2 weak derivative in the x_i-th direction one can use the fundamental theorem of calculus to prove that any bounded sequence in H is 2-equi-integrable.

3.2 Pure Bloch operators

Consider the space

$$\mathcal{V}_{\boldsymbol{\theta}} = \{v \in H^1_{\boldsymbol{\theta}}(Q) \mid \ v \equiv 0 \text{ on } Q_1\}, \tag{3.1}$$

which is a closed subspace of $H^1_{\boldsymbol{\theta}}(Q)$, and therefore is a Hilbert space when equipped with standard $H^1_{\boldsymbol{\theta}}(Q)$ norm. Define the sesquilinear form

$$\beta_{\boldsymbol{\theta}}(u,v) := \int_{Q_0} a_0(y)\boldsymbol{\nabla}_y u(y) \cdot \overline{\boldsymbol{\nabla}_y v}(y)\,\mathrm{d}y, \quad u, v \in \mathcal{V}_{\boldsymbol{\theta}}.$$

Since a_0 is positive and bounded on Q_0, and elements of $\mathcal{V}_{\boldsymbol{\theta}}$ have zero trace on the part of the boundary $\Gamma = \partial Q_1$, then by Poincaré's inequality the form $\beta_{\boldsymbol{\theta}}$ is (uniformly in $\boldsymbol{\theta}$) coercive and bounded on $\mathcal{V}_{\boldsymbol{\theta}}$, i.e. there exists c_1 and c_2 independent of $\boldsymbol{\theta}$ such that

$$|\beta_{\boldsymbol{\theta}}(u,v)| \le c_1 ||u||_{H^1_{\boldsymbol{\theta}}} ||v||_{H^1_{\boldsymbol{\theta}}},$$
$$\beta_{\boldsymbol{\theta}}(u,u) \ge c_2 ||u||^2_{H^1_{\boldsymbol{\theta}}},$$

for all $u, v \in \mathcal{V}_{\boldsymbol{\theta}}$. This implies that for every $f \in L^2(Q_0)$ there exists a unique solution $u \in \mathcal{V}_{\boldsymbol{\theta}}$ such that

$$\beta_{\boldsymbol{\theta}}(u,v) = \int_{Q_0} f(y)\overline{v}(y)\,\mathrm{d}y, \qquad \forall v \in \mathcal{V}_{\boldsymbol{\theta}}.$$

Consequently, the unbounded self-adjoint linear operator $B_{\boldsymbol{\theta}}$, defined in $L^2(Q_0)$, given by $B_{\boldsymbol{\theta}} u = f$, is positive and, moreover, by the Rellich embedding theorem has compact resolvent. Therefore the spectrum of $B_{\boldsymbol{\theta}}$ is discrete, and we order the eigenvalues in accordance with the min-max principle. These eigenvalues can be shown to be continuous functions of $\boldsymbol{\theta}$, in fact the following result holds.

Lemma 1. *For each $n \in \mathbb{N}$, let $\mu^{(n)}_{\boldsymbol{\theta}}$ denote the n-th eigenvalue of $B_{\boldsymbol{\theta}}$ as ordered according to the min-max principle, i.e.*

$$\mu^{(n)}_{\boldsymbol{\theta}} = \sup_{v_1,\dots,v_{n-1} \in \mathcal{V}_{\boldsymbol{\theta}}} \ \inf_{\substack{v \in \mathcal{V}_{\boldsymbol{\theta}}, \\ ||v||_{L^2(Q_0)}=1, \\ v \perp v_i, \forall i=1,\dots,n}} \int_{Q_0} a_0 \boldsymbol{\nabla} v \cdot \overline{\boldsymbol{\nabla} v}, \qquad \boldsymbol{\theta} \in [0,2\pi)^3, \tag{3.2}$$

where $v \perp v_i$ is shorthand for v is orthogonal to v_i in $L^2(Q_0)$. Then, for each $n \in \mathbb{N}$ the function $\lambda_n(\boldsymbol{\theta}) := \mu^{(n)}_{\boldsymbol{\theta}}$ is Lipschitz continuous, that is there exists a $C_n > 0$ such that

$$|\lambda_n(\boldsymbol{\theta}') - \lambda_n(\boldsymbol{\theta})| \le C_n |\boldsymbol{\theta}' - \boldsymbol{\theta}|, \qquad \forall \boldsymbol{\theta}, \boldsymbol{\theta}' \in [0,2\pi)^3.$$

The proof relies on an important observation is that the spaces $\mathcal{V}_{\boldsymbol{\theta}}$, $\boldsymbol{\theta} \in [0,2\pi)^3$, are mutually isomorphic. Indeed, if $\boldsymbol{\theta}, \boldsymbol{\theta}' \in [0,2\pi)^3$ then it is clear that the isometric mapping $\mathcal{U}(\boldsymbol{\theta}, \boldsymbol{\theta}') : L^2(Q) \to L^2(Q)$ defined as multiplication by the function $\exp\big(\mathrm{i}(\boldsymbol{\theta}' - \boldsymbol{\theta}) \cdot \mathrm{y}\big)$ defines an isomorphism between $\mathcal{V}_{\boldsymbol{\theta}}$ and $\mathcal{V}_{\boldsymbol{\theta}'}$.

Proof. Let v be $L^2(Q_0)$-normalised element of $\mathcal{V}_{\boldsymbol{\theta}}$ and consider $v' := \mathcal{U}(\boldsymbol{\theta}, \boldsymbol{\theta}')v = \exp\big(\mathrm{i}(\boldsymbol{\theta}' - \boldsymbol{\theta}) \cdot \mathrm{y}\big)\mathrm{v}$. Then, v' is an $L^2(Q_0)$-normalised element of $\mathcal{V}_{\boldsymbol{\theta}'}$ and the following identity

$$\int_{Q_0} a_0 \boldsymbol{\nabla} v' \cdot \overline{\boldsymbol{\nabla} v'} = \int_{Q_0} a_0(y)\boldsymbol{\nabla} v(y) \cdot \overline{\boldsymbol{\nabla} v}(y)\,\mathrm{d}y + \int_{Q_0} a_0(y)\boldsymbol{\nabla} v(y) \cdot \overline{\mathrm{i}(\boldsymbol{\theta}' - \boldsymbol{\theta})\mathrm{v}}(y)\,\mathrm{d}y$$
$$+ \int_{Q_0} a_0(y)\mathrm{i}(\boldsymbol{\theta}' - \boldsymbol{\theta})\exp\big(\mathrm{i}(\boldsymbol{\theta}' - \boldsymbol{\theta})\mathrm{y}\big)\mathrm{v}(\mathrm{y}) \cdot \overline{\boldsymbol{\nabla}\mathrm{v}'}(\mathrm{y})\,\mathrm{d}\mathrm{y}$$

holds. Therefore, one has

$$\left| \int_{Q_0} a_0 \boldsymbol{\nabla} v' \cdot \overline{\boldsymbol{\nabla} v'} - \int_{Q_0} a_0 \boldsymbol{\nabla} v \cdot \overline{\boldsymbol{\nabla} v} \right|$$
$$\le ||a_0||^{1/2}_{L^\infty(Q_0)} |\boldsymbol{\theta}' - \boldsymbol{\theta}| \left[\left(\int_{Q_0} a_0 \boldsymbol{\nabla} v \cdot \boldsymbol{\nabla} v \right)^{1/2} + \left(\int_{Q_0} a_0 \boldsymbol{\nabla} v' \cdot \boldsymbol{\nabla} v' \right)^{1/2} \right].$$

Consequently, as the isometric mapping $\mathcal{U}(\boldsymbol{\theta},\boldsymbol{\theta}') : L^2(Q_0) \to L^2(Q_0)$ is an isomorphism between $\mathcal{V}_{\boldsymbol{\theta}}$ and $\mathcal{V}_{\boldsymbol{\theta}'}$, the above inequality and the min-max formula (3.2) implies that

$$|\lambda_n(\boldsymbol{\theta}') - \lambda_n(\boldsymbol{\theta})| \leq ||a_0||^{1/2}_{L^\infty(Q_0)} |\boldsymbol{\theta}' - \boldsymbol{\theta}| \big(\lambda_n(\boldsymbol{\theta}') + \lambda_n(\boldsymbol{\theta})\big). \tag{3.3}$$

Now, if we consider the self-adjoint Dirichlet operator in $L^2(Q_0)$ associated with the form

$$\beta_D(u,v) := \int_{Q_0} a_0 \boldsymbol{\nabla} u \cdot \boldsymbol{\nabla} v, \qquad \forall u, v \in H^1_0(Q_0),$$

then, since $H^1_0(Q_0)$ is embedded in $\mathcal{V}_{\boldsymbol{\theta}}$ for all $\boldsymbol{\theta}$, one has

$$\lambda_n(\boldsymbol{\theta}) \leq \mu_n := \sup_{v_1,\dots,v_{n-1} \in H^1_0(Q_0)} \inf_{\substack{v \in H^1_0(Q_0), \\ ||v||_{L^2(Q_0)}=1, \\ v \perp v_i, \forall i=1,\dots,n}} \int_{Q_0} a_0 \boldsymbol{\nabla} v \cdot \boldsymbol{\nabla} v, \quad \forall \boldsymbol{\theta} \in [0,2\pi)^3.$$

Here μ_n is the n-th eigenvalue[4] of the operator B_D, defined in a similar manner as $B_{\boldsymbol{\theta}}$ above. Hence, we deduce from (3.3) that $\lambda_n(\boldsymbol{\theta})$ is Lipschitz continuous with a Lipschitz constant bounded from above by $2||a_0||^{1/2}_{L^\infty(Q_0)} \mu_n$. □

4 Limit spectrum

In this section we study the spectrum

$$\bigcup_{\boldsymbol{\theta} \in [0,2\pi)^3} \sigma(A^{\mathrm{hom}}_{\boldsymbol{\theta}}).$$

In particular we shall prove Proposition 1. Let us fix $\boldsymbol{\theta} \in [0,2\pi)^3$ and suppose that $\lambda_{\boldsymbol{\theta}}$ is in the spectrum of $A^{\mathrm{hom}}_{\boldsymbol{\theta}}$. Then, there exists an eigenfunction $u_{\boldsymbol{\theta}} \in V_{\boldsymbol{\theta}}$ that solves the spectral problem

$$\begin{cases} -\mathrm{div}_y\big(a_0(y)\boldsymbol{\nabla}_y u_{\boldsymbol{\theta}}(x,y)\big) = \lambda_{\boldsymbol{\theta}} u_{\boldsymbol{\theta}}(x,y), & x \in \Omega,\ y \in Q_0, \\ \quad u_{\boldsymbol{\theta}}(x,y) = u_i(x), & x \in \Omega,\ y \in \overline{C_i}, \\ \text{where } u_i \equiv 0 \text{ if } \theta_i \neq 0 \text{ or otherwise solves} & \\ -a^{\mathrm{hom}}_i \partial^2_{x_i} u_i(x) + \mathcal{T}_i(u_{\boldsymbol{\theta}})(x) = \lambda |C_i| u_i(x), & \text{for } x \in \Omega. \end{cases} \tag{4.1}$$

Here, we recall that

$$\mathcal{T}_i(u_{\boldsymbol{\theta}})(x) = \int_{\Gamma_i} a_0(y) \boldsymbol{\nabla}_y u_{\boldsymbol{\theta}}(x,y) \cdot \mathbf{n}(y)\, \mathrm{d}S(y).$$

There are two sub cases to study, when $\boldsymbol{\theta} \in \cup_{i \in \mathcal{I}} H_i$, for $H_i := \{\boldsymbol{\theta} \in [0,2\pi^2)^3 \,|\, \boldsymbol{\theta}_i = 0\}$, and $\boldsymbol{\theta} \in [0,2\pi)^3 \backslash \big(\cup_{i \in \mathcal{I}} H_i\big)$.

Pure Bloch spectrum. If $\boldsymbol{\theta} \in [0,2\pi)^3 \backslash \big(\cup_{i \in \mathcal{I}} H_i\big)$, then $\lambda_{\boldsymbol{\theta}}, u_{\boldsymbol{\theta}}$ solves the problem

$$\begin{cases} -\mathrm{div}_y\big(a_0(y)\boldsymbol{\nabla}_y u_{\boldsymbol{\theta}}(x,y)\big) = \lambda_{\boldsymbol{\theta}} u_{\boldsymbol{\theta}}(x,y), & x \in \Omega,\ y \in Q_0, \\ \quad u_{\boldsymbol{\theta}}(x,y) = 0, & x \in \Omega,\ y \in \Gamma. \end{cases} \tag{4.2}$$

Therefore, setting $u_{\boldsymbol{\theta}}(x,y) = \phi(x) v_{\boldsymbol{\theta}}(y)$ for a sufficiently arbitrary ϕ, we find that $v_{\boldsymbol{\theta}}$ solves

$$\begin{cases} -\mathrm{div}_y\big(a_0(y)\boldsymbol{\nabla}_y v_{\boldsymbol{\theta}}(y)\big) = \lambda_{\boldsymbol{\theta}} v_{\boldsymbol{\theta}}(y), & y \in Q_0, \\ \quad v_{\boldsymbol{\theta}}(y) = 0, & y \in \Gamma. \end{cases} \tag{4.3}$$

Therefore, the spectrum of $A^{\mathrm{hom}}_{\boldsymbol{\theta}}$ for $\boldsymbol{\theta} \in [0,2\pi)^3 \backslash \big(\cup_{i \in \mathcal{I}} H_i\big)$ consists of eigenvalues of infinity multiplicity, and these eigenvalues coincide with the eigenvalues of the pure Bloch operator $B_{\boldsymbol{\theta}}$ associated to problem

[4]The spectrum of B_D is discrete, which again is a consequence of the Rellich theorem.

(4.3), see Section 3.2. Therein, the eigenvalues of the $B_{\boldsymbol{\theta}}$ are found to be continuous with respect to $\boldsymbol{\theta}$, and by continuously extending $\boldsymbol{\theta}$ from $[0,2\pi)^3\backslash\big(\cup_{i\in\mathcal{I}} H_i\big)$ to $[0,2\pi)^3$ we deduce that

$$\sigma(A^{\rm hom}) \supset \bigcup_{\boldsymbol{\theta}\in[0,2\pi)^3} \sigma(B_{\boldsymbol{\theta}}).$$

It is for this reason that we call $\bigcup_{\boldsymbol{\theta}\in[0,2\pi)^3}\sigma(B_{\boldsymbol{\theta}})$ the pure Bloch spectrum of $A^{\rm hom}_{\boldsymbol{\theta}}$.

Spatial spectrum. Let us now suppose that $\boldsymbol{\theta}\in\cup_{i\in\mathcal{I}}H_i$ and $\lambda_{\boldsymbol{\theta}}\in\sigma(A_{\boldsymbol{\theta}})$ is not a pure Bloch eigenvalue, i.e. $\lambda_{\boldsymbol{\theta}}\notin\bigcup_{\boldsymbol{\theta}\in[0,2\pi)^3}\sigma(B_{\boldsymbol{\theta}})$. Introducing, for $i\in\mathcal{I}_{\boldsymbol{\theta}}$ the functions $b^{(i)}_{\boldsymbol{\theta}}\in H^1_{\boldsymbol{\theta}}(Q)$ that satisfy

$$\begin{cases} -\mathrm{div}_y\big(a_0(y)\boldsymbol{\nabla}_y b^{(i)}_{\boldsymbol{\theta}}(y)\big)=0, & y\in Q_0,\\ b^{(i)}_{\boldsymbol{\theta}}(y)=\delta_{ij}, & y\in\overline{C_j},\ j=1,2,3,\end{cases} \tag{4.4}$$

we represent $u_{\boldsymbol{\theta}}$ as follows

$$u_{\boldsymbol{\theta}}(x,y)=\sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}}u_i(x)b^{(i)}_{\boldsymbol{\theta}}(y)+v_{\boldsymbol{\theta}}(x,y),$$

and substitute this representation into (4.1) to deduce that $v_{\boldsymbol{\theta}}(x,y)\in\mathcal{V}_{\boldsymbol{\theta}}$, see (3.1), solves

$$-\mathrm{div}_y\big(a_0(y)\boldsymbol{\nabla}_y v_{\boldsymbol{\theta}}(x,y)\big)-\lambda_{\boldsymbol{\theta}}v_{\boldsymbol{\theta}}(x,y)=\lambda_{\boldsymbol{\theta}}\sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}}u_i(x)b^{(i)}_{\boldsymbol{\theta}}(y),\quad x\in\Omega,y\in Q_0. \tag{4.5}$$

Denoting respectively by $\mu^{(m)}_{\boldsymbol{\theta}}$ and $v^{(m)}_{\boldsymbol{\theta}}$ the m-th eigenvalue and orthonormal eigenfunction of $B_{\boldsymbol{\theta}}$, we perform a spectral decomposition of $v_{\boldsymbol{\theta}}$ and $b^{(i)}_{\boldsymbol{\theta}}$ to conclude that

$$v_{\boldsymbol{\theta}}(x,y)=\sum_{m\in\mathbb{N}}c_m(\boldsymbol{\theta},x)v^{(m)}_{\boldsymbol{\theta}}(y),\quad b^{(i)}_{\boldsymbol{\theta}}(y)=\sum_{m\in\mathbb{N}}b^{(i)}_m(\boldsymbol{\theta})v^{(m)}_{\boldsymbol{\theta}}(y),$$

for some $c_m(\boldsymbol{\theta},x)$ and constants $b^{(i)}_m(\boldsymbol{\theta})=\int_{Q_0}b^{(i)}_{\boldsymbol{\theta}}\overline{v^{(m)}_{\boldsymbol{\theta}}}$. Substituting the spectral representations into (4.5) gives

$$c_m(\boldsymbol{\theta},x)=\tfrac{\lambda_{\boldsymbol{\theta}}}{\mu^{(m)}_{\boldsymbol{\theta}}-\lambda_{\boldsymbol{\theta}}}\sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}}u_i(x)b^{(i)}_m(\boldsymbol{\theta}).$$

Therefore, $u_{\boldsymbol{\theta}}$ admits the form

$$u_{\boldsymbol{\theta}}(x,y)=\sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}}\sum_{m\in\mathbb{N}}\Big(\tfrac{\mu^{(m)}_{\boldsymbol{\theta}}}{\mu^{(m)}_{\boldsymbol{\theta}}-\lambda_{\boldsymbol{\theta}}}\Big)u_i(x)b^{(i)}_m(\boldsymbol{\theta})v^{(m)}_{\boldsymbol{\theta}}(y),\qquad x\in\Omega,y\in Q_0.$$

Consequently, we calculate

$$\mathcal{T}_j(u_{\boldsymbol{\theta}})(x)=\sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}}\sum_{m\in\mathbb{N}}\Big(\tfrac{\mu^{(m)}_{\boldsymbol{\theta}}}{\mu^{(m)}_{\boldsymbol{\theta}}-\lambda_{\boldsymbol{\theta}}}\Big)u_i(x)b^{(i)}_m(\boldsymbol{\theta})\int_{\Gamma_j}a_0\boldsymbol{\nabla}_y v^{(m)}_{\boldsymbol{\theta}}(y)\cdot\mathbf{n}(y)\,\mathrm{d}S(y).$$

Recalling that $b^{(i)}_{\boldsymbol{\theta}}$ solves (4.4), $v^{(m)}_{\boldsymbol{\theta}}$ solves (4.3), and utilising Green's identity, we deduce that

$$\begin{aligned}\int_{\Gamma_j}a_0\boldsymbol{\nabla}_y v^{(m)}_{\boldsymbol{\theta}}(y)\cdot\mathbf{n}(y)\,\mathrm{d}S(y)&=\int_{\Gamma}a_0\boldsymbol{\nabla}_y v^{(m)}_{\boldsymbol{\theta}}(y)\cdot\mathbf{n}(y)\overline{b^{(j)}_{\boldsymbol{\theta}}}(y)\,\mathrm{d}S(y)=-\mu^{(m)}_{\boldsymbol{\theta}}\int_{Q_0}v^{(m)}_{\boldsymbol{\theta}}\overline{b^{(j)}_{\boldsymbol{\theta}}}\\&=-\mu^{(m)}_{\boldsymbol{\theta}}\overline{b^{(j)}_m}(\boldsymbol{\theta}).\end{aligned}$$

Therefore

$$\mathcal{T}_j(u_{\boldsymbol{\theta}})(x)=-\sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}}\sum_{m\in\mathbb{N}}\Big(\tfrac{|\mu^{(m)}_{\boldsymbol{\theta}}|^2}{\mu^{(m)}_{\boldsymbol{\theta}}-\lambda_{\boldsymbol{\theta}}}\Big)u_i(x)b^{(i)}_m(\boldsymbol{\theta})\overline{b^{(j)}_m(\boldsymbol{\theta})},$$

and for each $i \in \mathcal{I}_{\boldsymbol{\theta}}$, u_i solves the problem

$$-a_i^{\hom}\partial_{x_i}^2 u_i(x) = \sum_{j\in\mathcal{I}_{\boldsymbol{\theta}}} \beta_{\boldsymbol{\theta}}^{(ij)}(\lambda_{\boldsymbol{\theta}})u_j(x) \qquad x\in\Omega, \qquad u_i\nu_i = 0 \text{ on } \partial\Omega,$$

for

$$\beta_{\boldsymbol{\theta}}^{(ij)}(\lambda) = \lambda|C_i|\delta_{ij} + \sum_{m\in\mathbb{N}} \Big(\tfrac{|\mu_{\boldsymbol{\theta}}^{(m)}|^2}{\mu_{\boldsymbol{\theta}}^{(m)}-\lambda}\Big) b_m^{(j)}(\boldsymbol{\theta})\overline{b_m^{(i)}(\boldsymbol{\theta})}, \qquad \lambda\in\mathbb{R}. \tag{4.6}$$

Hence, we have demonstrated Proposition 1.

5 Quasi-periodic two-scale convergence

In this section we introduce an appropriate notion of convergence that will account for the presence of microscopic oscillations that are quasi-periodic in nature. This convergence will turn out to be a natural extension of the standard (periodic) two-scale convergence introduced by G. Nguetseng [25]-G. Allaire [1]. In particular, we aim to use this extended notion of two-scale convergence to study the spectral properties of operator families in homogenisation theory in a similar vein to that first introduced by V. V. Zhikov [28,29]. Throughout this section Ω is a domain in $\mathbb{R}^d$, $d\geq 1$, $\square := (0,1)^d$ and $\boldsymbol{\theta}\in[0,2\pi)^d$. All of the functions, even if real-valued, are considered to take values in the complex field. The space $C_{\boldsymbol{\theta}}^\infty(\square)$ shall denote the space of smooth functions $\varphi(y)$ whose functions and derivatives are $\boldsymbol{\theta}$-quasi-periodic with respect to y: $\varphi(y+\mathbf{e}_j) = \exp(\mathrm{i}\boldsymbol{\theta}_\mathrm{j})\varphi(\mathrm{y})$ for each $y\in\square$ and each Euclidean basis vector $\mathbf{e}_j$, $j=1,\dots,d$. Note, that for elements in this space the following mean-value property holds:

Mean-value property. For every $\varphi\in C_{\boldsymbol{\theta}}^\infty(\square)$ the following convergence holds:

$$\lim_{\varepsilon\to 0}\int_\Omega \left|\phi(x)\varphi\left(\tfrac{x}{\varepsilon}\right)\right|^2 \,\mathrm{d}x = \int_\Omega\int_\square |\phi(x)\varphi(y)|^2, \quad \forall\phi\in C_0^\infty(\Omega). \tag{5.1}$$

This fact follows by noting that the property holds for elements in $C_\#^\infty(\square)$ the space of smooth periodic functions , see for example [1, Lemma 1.3], and observing that multiplication by $\exp(-\mathrm{i}\boldsymbol{\theta}\cdot\mathrm{y})$ defines an isomorphism between $C_{\boldsymbol{\theta}}^\infty(\square)$ and $C_\#^\infty(\square)$ that preserves absolute value. Indeed, φ belongs to $C_{\boldsymbol{\theta}}^\infty(\square)$ if, and only if, $\exp(-\mathrm{i}\boldsymbol{\theta}\cdot\mathrm{y})\varphi$ belongs to $C_\#^\infty(\square)$ and $|\varphi|(y) = |\exp(-\mathrm{i}\boldsymbol{\theta}\cdot\mathrm{y})\varphi|(\mathrm{y})$ in $\square$.

The remainder of this section is dedicated to the introduction of the notion $\boldsymbol{\theta}$-quasi-periodic two-scale convergence and an exposition of results that are used throughout the article to establish the main result. We remark here that because $C_{\boldsymbol{\theta}}^\infty(\square)$ is isomorphic to $C_\#^\infty(\square) = C_{\mathbf{0}}^\infty(\square)$, with isomorphism $\exp\big(-\mathrm{i}\boldsymbol{\theta}\cdot\mathrm{y}\big)$, then the results presented in this section[5] are immediately established for each $\boldsymbol{\theta}\in[0,2\pi)^3$ if proved for $\boldsymbol{\theta}=\mathbf{0}$. We shall demonstrate this with the first result of the section and omit the remaining proofs which follow in a similar vein.

Definition 1. Let $u_\varepsilon\in L^2(\Omega)$ be a bounded sequence and $u\in L^2(\Omega\times\square)$. Then, we say u_ε (weakly) $\boldsymbol{\theta}$-quasi-periodic two-scale converges to u, denoted by $u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} u$, if the following convergence

$$\lim_{\varepsilon\to 0}\int_\Omega u_\varepsilon(x)\overline{\phi(x)\varphi\left(\tfrac{x}{\varepsilon}\right)}\,\mathrm{d}x = \int_\Omega\int_\square u(x,y)\overline{\phi(x)\varphi(y)}\,\mathrm{d}y\mathrm{d}x, \qquad \forall\phi\in C_0^\infty(\Omega),\ \forall\varphi\in C_{\boldsymbol{\theta}}^\infty(\square) \tag{5.2}$$

holds.

Remark 3. Notice that for $\boldsymbol{\theta}=0$, this is the standard notion of two-scale convergence.

The next important result states that bounded sequences in $L^2(\Omega)$ are relatively compact with respect to $\boldsymbol{\theta}$-quasi-periodic two-scale convergence.

[5]The results in this section can be established by first principles making no reference to such an isomorphism.

Proposition 2. *If u_ε is bounded in $L^2(\Omega)$ then, up to a subsequence, u_ε weakly $\boldsymbol{\theta}$-quasi-periodic two-scale converges to some $u \in L^2(\Omega \times \square)$.*

Proof. The result has been established previously for the case $\boldsymbol{\theta} = \mathbf{0}$, see for example [1, 25, 28]. Let us consider $\boldsymbol{\theta} \neq \mathbf{0}$. Note that the function $\widetilde{u}_\varepsilon = \exp\big(-\mathrm{i}\boldsymbol{\theta}\cdot \mathrm{y}\big)\mathrm{u}_\varepsilon$ is bounded in L^2 and therefore by the assertion for $\boldsymbol{\theta} = \mathbf{0}$, up to a discarded subsequence, $\widetilde{u}_\varepsilon$ ($\mathbf{0}$-quasi-periodically) two-scale converges to some $\widetilde{u} \in L^2(\Omega \times Q)$. Now, the result follows from this fact and noting that for fixed $\varphi \in C^\infty_{\boldsymbol{\theta}}(\square)$ one has

$$\int_\Omega u_\varepsilon(x)\overline{\phi(x)\varphi(\tfrac{x}{\varepsilon})}\,\mathrm{d}x = \int_\Omega \widetilde{u}_\varepsilon(x)\overline{\phi(x)\exp\big(-\mathrm{i}\boldsymbol{\theta}\cdot\tfrac{\mathrm{x}}{\varepsilon}\big)\varphi(\tfrac{\mathrm{x}}{\varepsilon})}\,\mathrm{d}x,$$
$$\int_\Omega\int_\square \exp\big(-\mathrm{i}\boldsymbol{\theta}\cdot\mathrm{y}\big)\widetilde{\mathrm{u}}(\mathrm{x},\mathrm{y})\overline{\phi(\mathrm{x})\varphi(\mathrm{y})}\,\mathrm{dydx} = \int_\Omega\int_{\mathrm{Q}} \widetilde{\mathrm{u}}(\mathrm{x},\mathrm{y})\overline{\phi(\mathrm{x})\exp\big(-\mathrm{i}\boldsymbol{\theta}\cdot\mathrm{y}\big)\varphi(\mathrm{y})}\,\mathrm{dydx},$$

and that $\exp\big(-\mathrm{i}\boldsymbol{\theta}\cdot\mathrm{y}\big)\varphi$ is a smooth periodic function. Hence $u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} \exp\big(-\mathrm{i}\boldsymbol{\theta}\cdot\mathrm{y}\big)\widetilde{\mathrm{u}}$. □

An important result from the point of view of homogenisation theory is that the test functions φ in (5.2) can be taken to be $\boldsymbol{\theta}$-quasi-periodic elements of $L^2(\square)$, i.e. the following result holds.

Proposition 3. *If $u_\varepsilon \in L^2(\Omega)$ $\boldsymbol{\theta}$-quasi-periodic two-scale converges to $u \in L^2(\Omega \times \square)$, then the following convergence*

$$\lim_{\varepsilon\to 0}\int_\Omega u_\varepsilon(x)\overline{\phi(x)\psi\left(\tfrac{x}{\varepsilon}\right)}\,\mathrm{d}x = \int_\Omega\int_\square u(x,y)\overline{\phi(x)\psi(y)}\,\mathrm{d}y\mathrm{d}x$$

holds for all $\phi \in C_0^\infty(\Omega)$, and for all $\psi \in L^2(\square)$ such that $\psi(y+\mathbf{e}_j) = \exp(\mathrm{i}\boldsymbol{\theta}_\mathrm{j})\psi(\mathrm{y})$ for almost every $y \in \square$ and $j = 1,\dots,d$.

Remark 4. If Ω is a bounded domain, as in this article, then additionally the test functions ϕ can be taken to be elements of $C(\overline{\Omega})$.

The following results are of interest, but will not be directly used in the article.

Proposition 4.

1. *For $u_\varepsilon \in L^2(\Omega)$ $\boldsymbol{\theta}$-quasi-periodic two-scale converging to $u \in L^2(\Omega \times \square)$ one has that*

$$\exp(-\mathrm{i}\boldsymbol{\theta}\cdot\tfrac{\mathrm{x}}{\varepsilon})\mathrm{u}_\varepsilon(\mathrm{x}) \rightharpoonup \int_\square \exp(-\mathrm{i}\boldsymbol{\theta}\cdot\mathrm{y})\mathrm{u}(\mathrm{x},\mathrm{y})\,\mathrm{dy} \quad \textit{weakly in } L^2(\Omega).$$

2. *For $u_\varepsilon \in L^2(\Omega)$ $\boldsymbol{\theta}$-quasi-periodic two-scale converging to $u \in L^2(\Omega \times \square)$ then*

$$\liminf_{\varepsilon\to 0}\int_\Omega |u_\varepsilon(x)|^2\,\mathrm{d}x \geq \int_\Omega\int_\square |u(x,y)|^2\,\mathrm{d}y\mathrm{d}x.$$

A result of particular interest in high-contrast homogenisation problems is as follows.

Proposition 5. *Let $u_\varepsilon \in H^1(\Omega)$ satisfy*

$$\sup_\varepsilon ||u_\varepsilon||_{L^2(\Omega)} < \infty, \qquad \sup_\varepsilon ||\varepsilon\boldsymbol{\nabla} u_\varepsilon||_{L^2(\Omega)} < \infty.$$

Then, there exists $u \in L^2(\Omega; H^1_{\boldsymbol{\theta}}(\square))$ such that, up to a subsequence, the following convergences hold:

$$u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} u, \qquad \varepsilon\boldsymbol{\nabla} u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} \boldsymbol{\nabla}_y u.$$

Proof. Let ϕ and $\mathbf{\Psi}$ denote respectively fixed arbitrary elements of $C_0^\infty(\Omega)$ and $C_{\boldsymbol{\theta}}^\infty(\square;\mathbb{C}^d)$. By Proposition 2, there exists $u \in L^2(\Omega\times\square)$ and $\boldsymbol{\chi} \in L^2(\Omega\times\square;\mathbb{C}^d)$ such that, up to a subsequence that we discard, the following convergences hold:

$$u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} u, \qquad \varepsilon\boldsymbol{\nabla} u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} \chi. \tag{5.3}$$

To prove that $u \in L^2(\Omega; H^1_{\boldsymbol{\theta}}(Q))$, we pass to the limit in the identity

$$\begin{aligned}\int_\Omega \varepsilon\boldsymbol{\nabla} u_\varepsilon(x)\cdot\overline{\phi(x)\mathbf{\Psi}(\tfrac{x}{\varepsilon})}\,\mathrm{d}x &= -\int_\Omega u_\varepsilon(x)\overline{\varepsilon\mathrm{div}\big(\phi(x)\mathbf{\Psi}(\tfrac{x}{\varepsilon})\big)}\,\mathrm{d}x\\ &= -\int_\Omega u_\varepsilon(x)\overline{\varepsilon\boldsymbol{\nabla}_x\phi(x)\cdot\mathbf{\Psi}(\tfrac{x}{\varepsilon})}\,\mathrm{d}x - \int_\Omega u_\varepsilon(x)\overline{\phi(x)\mathrm{div}_y\mathbf{\Psi}(\tfrac{x}{\varepsilon})}\,\mathrm{d}x,\end{aligned}$$

using, the Mean-value property (5.1) for $\mathbf{\Psi}$, the boundedness of u_ε, and (5.3) to deduce that

$$\int_\Omega\int_\square \boldsymbol{\chi}(x,y)\cdot\overline{\phi(x)\mathbf{\Psi}(y)}\,\mathrm{d}y\mathrm{d}x = -\int_\Omega\int_\square u(x,y)\overline{\phi(x)\mathrm{div}_y\mathbf{\Psi}(y)}\,\mathrm{d}y\mathrm{d}x.$$

Therefore, for almost every x, the functions $\boldsymbol{\chi}(x,\cdot)$ and $u(x,\cdot)$ are related by the identity

$$\int_\square \boldsymbol{\chi}(x,y)\cdot\overline{\mathbf{\Psi}(y)}\,\mathrm{d}y = -\int_\square u(x,y)\overline{\mathrm{div}_y\mathbf{\Psi}(y)}\,\mathrm{d}y, \qquad \forall\mathbf{\Psi}\in C_{\boldsymbol{\theta}}^\infty(\square;\mathbb{C}^d),$$

that is $u \in L^2(\Omega; H^1_{\boldsymbol{\theta}}(\square))$ and $\boldsymbol{\nabla}_y u = \boldsymbol{\chi}$. □

We end this section with a result that is illuminating when it comes to studying the convergence of spectra for parameter-dependent operator families. It gives a one-sided justification for the convergence of the spectra associated to high-contrast problems to the spectrum associated to the $\boldsymbol{\theta}$-quasi-periodic two-scale limit problems.

Definition 2. Fix $\boldsymbol{\theta}\in[0,2\pi)^d$, and let A_ε and A be non-negative self-adjoint operators defined in $L^2(\Omega)$ and H a closed subset of $L^2(\Omega\times\square)$. Then, we say that A_ε strong $\boldsymbol{\theta}$-quasi-periodic two-scale resolvent converges to A if for every $f_\varepsilon(x)\in L^2(\Omega)$ that $\boldsymbol{\theta}$-quasi-periodic two-scale converge to $f(x,y)\in L^2(\Omega\times\square)$, the following convergence

$$u_\varepsilon = (A_\varepsilon + I)^{-1}f_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} u = (A+I)^{-1}Pf, \text{ as } \varepsilon\to 0$$

holds. Here, P is the orthogonal projection onto H in $L^2(\Omega\times\square)$.

Here, we state an important consequence of such resolvent convergence. The proof, omitted here, follows standard spectral theoretic arguments, see for example [28].

Proposition 6. *If A_ε strong $\boldsymbol{\theta}$-quasi-periodic two-scale resolvent converges to A then the spectrum $\sigma(A)$ of A is related to the spectrum $\sigma(A_\varepsilon)$ of A_ε in the following sense.*

For every $\lambda\in\sigma(A)$ there exists $\lambda_\varepsilon\in\sigma(A_\varepsilon)$ such that λ_ε converges to λ as ε tends to zero.

6 Proof of the homogenisation theorem

This section is dedicated to the proof of Theorem 1. To do this, we shall develop an appropriate quasi-periodic two-scale variation of a powerful method first introduced in [20] for the case of standard (periodic) two-scale convergence, i.e. $\boldsymbol{\theta}$-quasi-periodic two-scale convergence for $\boldsymbol{\theta}=0$. In what follows ϕ, φ and $\mathbf{\Phi}$ will denote respectively fixed arbitrary elements of $C_0^\infty(\Omega)$, $C_{\boldsymbol{\theta}}^\infty(Q)$ and $C_{\boldsymbol{\theta}}^\infty(Q;\mathbb{C}^3)$.

6.1 Technical preliminaries

The following results will be of importance in the proof of the homogenisation theorem.

Lemma 2. *Let B be the closure of a smooth domain and let B_1 be a smooth bounded domain such that $B \subset B_1$ and $A = B_1 \backslash B$ is a connected set. Then every $u \in H^1\big((0,1)\times A\big)$ can be extended to $(0,1)\times B_1$ as a function $\widetilde{u} \in H^1\big((0,1)\times B_1\big)$ such that*

$$\begin{aligned} \int_{(0,1)\times B_1} |\boldsymbol{\nabla}\widetilde{u}|^2 &\leq c \int_{(0,1)\times A} |\boldsymbol{\nabla} u|^2, \\ \int_{(0,1)\times B_1} |\widetilde{u}|^2 &\leq c \int_{(0,1)\times A} |u|^2, \end{aligned} \tag{6.1}$$

where c does not depend on $u \in H^1((0,1)\times A)$.

Proof. Suppose $u \in H^1\big((0,1)\times A\big)$. Then, by Fubini's theorem, for almost every $x_1 \in (0,1)$ the function $u(x_1,\cdot)$ belongs to $H^1(A)$ and let $Eu(x_1,\cdot)$ be the Sobolev extension of $u(x_1,\cdot)$ into B_1 given in [18, Lemma 3.2, pg. 88]. In particular, one has

$$\begin{aligned} \int_{B_1} |\boldsymbol{\nabla}' Eu(x_1,\cdot)|^2 &\leq c \int_A |\boldsymbol{\nabla}' u(x_1,\cdot)|^2, \\ \int_{B_1} |Eu(x_1,\cdot)|^2 &\leq c \int_A |u(x_1,\cdot)|^2, \end{aligned} \tag{6.2}$$

where c does not depend on u nor x_1. Here, $\boldsymbol{\nabla}'$ denotes the gradient vector $(0, \partial_{x_2}, \partial_{x_3})$.
Consider $\widetilde{u}$ given by

$$\widetilde{u(x_1,\cdot)} := Eu(x_1,\cdot), \qquad \text{a.e. } x_1 \in (0,1). \tag{6.3}$$

Then $\boldsymbol{\nabla}'\widetilde{u(x_1,\cdot)} = \boldsymbol{\nabla}' Eu(x_1,\cdot)$ and from assertion (6.2) it follows that

$$\begin{aligned} \int_{(0,1)\times B_1} |\boldsymbol{\nabla}'\widetilde{u}|^2 &\leq c \int_{(0,1)\times A} |\boldsymbol{\nabla}' u|^2, \\ \int_{(0,1)\times B_1} |\widetilde{u}|^2 &\leq c \int_{(0,1)\times A} |u|^2. \end{aligned}$$

To prove (6.1), it remains to demonstrate that $\partial_{x_1}\widetilde{u} \in L^2\big((0,1)\times B_1\big)$ and

$$\int_0^1 \int_{B_1} |\partial_{x_1}\widetilde{u}|^2(x_1,x')\,\mathrm{d}x'\mathrm{d}x_1 \leq c \int_0^1 \int_A |\partial_{x_1} u|^2(x_1,x')\,\mathrm{d}x'\mathrm{d}x_1. \tag{6.4}$$

For each $t \in \mathbb{R}$, the difference quotient is given by

$$D_t\widetilde{u}(x_1,x') := \frac{\widetilde{u}(x_1+t,x') - \widetilde{u}(x_1,x')}{t},$$

where we have extended $\widetilde{u}$ trivially by zero into $\mathbb{R}\backslash(0,1)$. Notice that $D_t\widetilde{u} = ED_t u$, i.e. the extension into B_1 of the function $\frac{u(x_1+t,\cdot)-u(x_1,\cdot)}{t} = D_t u(x_1+t,\cdot)$, and consequently

$$\int_0^1 \int_{B_1} |D_t\widetilde{u}|^2(x_1,x')\,\mathrm{d}x'\mathrm{d}x_1 \leq c \int_0^1 \int_A |D_t u|^2(x_1,x')\,\mathrm{d}x'\mathrm{d}x_1.$$

Since $D_t u$ converges strongly in $L^2\big((0,1)\times A\big)$ to $\partial_{x_1} u$, it follows that $D_t\widetilde{u}$ is a Cauchy sequence in $L^2\big((0,1)\times B_1\big)$ and this limit can be identified, using the fact that $(D_t\widetilde{u},\phi)_{L^2} = (\widetilde{u}, D_{-t}\phi)_{L^2}$ for $\phi \in C_0^\infty\big((0,1)\times B_1\big)$, as $\partial_{x_1}\widetilde{u}$. Furthermore, passing to the limit in the above inequality yields (6.4). □

Proposition 7. *Fix $\boldsymbol{\theta} \in [0, 2\pi)^3$. There exists a constant $C = C_{\boldsymbol{\theta}} > 0$ such that*

$$\inf_{v \in V_{\boldsymbol{\theta}}} ||u - v||_{H^1_{\boldsymbol{\theta}}(Q)} \leq C_{\boldsymbol{\theta}} ||\sqrt{a_1} \boldsymbol{\nabla} u||_{L^2(Q)}, \qquad \forall u \in H^1_{\boldsymbol{\theta}}(Q).$$

Here $V_{\boldsymbol{\theta}} = \ker \sqrt{a_1} \boldsymbol{\nabla}_{\boldsymbol{\theta}} = \{u \in H^1_{\boldsymbol{\theta}}(Q) \,\big|\, \sqrt{a_1} \boldsymbol{\nabla} u \equiv 0\}$.

Proof. For each $i \in \mathcal{I}$, let S_i be the cross-section of the cylinder C_i. Since S_i are compactly contained in $(0,1)^2$ and have mutually disjoint closures then there exists open A_i such that $\overline{S_i} \subset A_i \subset (0,1)^2$ and A_i are mutually disjoint. Let $\chi_i \in C_0^\infty(A_i)$ be smooth cut-off functions that are identity on S_i, we extend χ_i by zero to $(0,1)^2$.
Now using Lemma 2, let $\widetilde{u_i}$ be the extension of $u|_{C_i} \in H^1(C_i)$ to $H^1(D_i)$, where D_i is the cylinder whose axis is parallel to y_i with cross-section A_i. Note that since u is θ_i-quasi-periodic in the variable y_i, then the extension will be also, see (6.3). By Lemma 2 it follows that

$$\int_{D_i} |\boldsymbol{\nabla} \widetilde{u_i}|^2 \leq c \int_{C_i} |\boldsymbol{\nabla} u_i|^2. \tag{6.5}$$

For $\theta_i \neq 0$, the following Poincaré inequality

$$\int_{D_i} |\widetilde{u_i}|^2 \leq |\theta_i|^{-2} \int_{D_i} |\boldsymbol{\nabla} \widetilde{u_i}|^2 \tag{6.6}$$

holds[6]. For $\theta_i = 0$, one has

$$\int_{D_i} |\widetilde{u_i} - \langle \widetilde{u_i} \rangle|^2 \leq C \int_{D_i} |\boldsymbol{\nabla} \widetilde{u_i}|^2 \tag{6.7}$$

for some $C > 0$. Here $\langle \widetilde{u_i} \rangle := \frac{1}{D_i} \int_{D_i} \widetilde{u_i}$.
Recalling, $\mathcal{I}_{\boldsymbol{\theta}} = \{i \in \mathcal{I} \,|\, \theta_i = 0\}$, we set $\widetilde{u} = \sum_{i \in \mathcal{I}_{\boldsymbol{\theta}}} \chi_i (\widetilde{u_i} - \langle \widetilde{u_i} \rangle) + \sum_{i \in \mathcal{I} \setminus \mathcal{I}_{\boldsymbol{\theta}}} \chi_i \widetilde{u_i}$, here χ_i are taken to be constant in the variable y_i and as above the complementary directions. It follows that $\widetilde{u} \in H^1_{\boldsymbol{\theta}}(Q)$ and $u - \widetilde{u} \in V_{\boldsymbol{\theta}}$. Note that, by construction and inequalities (6.6), and (6.7), one has

$$||\widetilde{u}||^2_{H^1(Q)} \leq c_{\boldsymbol{\theta}} \sum_{i \in \mathcal{I}} ||\boldsymbol{\nabla} \widetilde{u_i}||^2_{L^2(D_i)}.$$

Now, the positivity of a_1 on Q_1 and (6.5) imply that the element $v := u - \widetilde{u}$ of $V_{\boldsymbol{\theta}}$ is such that

$$||u - v||^2_{H^1(Q)} = ||\widetilde{u}||^2_{H^1(Q)} \leq C_{\boldsymbol{\theta}} \int_{Q_1} a_1 |\boldsymbol{\nabla} u|^2,$$

and the result follows.

□

6.2 Proof of Theorem 1

Consider the sequence u_ε of solutions to (2.3), i.e.

$$\int_\Omega \big(a_1(\tfrac{x}{\varepsilon}) + \varepsilon^2 a_0(\tfrac{x}{\varepsilon})\big) \boldsymbol{\nabla} u_\varepsilon(x) \cdot \overline{\boldsymbol{\nabla} \phi}(x) \,\mathrm{d}x \; + \int_\Omega u_\varepsilon(x) \overline{\phi}(x) \,\mathrm{d}x = \int_\Omega f_\varepsilon(x) \overline{\phi}(x) \,\mathrm{d}x, \qquad \forall \phi \in C_0^\infty(\Omega). \tag{6.8}$$

for $f_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} f$, and recall, cf. (2.4), that

$$\sup_\varepsilon \big(||\sqrt{a_1}(\tfrac{\cdot}{\varepsilon}) \boldsymbol{\nabla} u_\varepsilon||_{L^2(\Omega;\mathbb{C}^3)} + ||\varepsilon \boldsymbol{\nabla} u_\varepsilon||_{L^2(\Omega;\mathbb{C}^3)} + \;\; ||u_\varepsilon||_{L^2(\Omega)} \big) < \infty. \tag{6.9}$$

[6]This follows from noting the lower bound on the spectrum of the laplacian on the space of $H^1\boldsymbol{\theta}(D_i)$ functions that are θ_i-quasi-periodic in direction y_i.

Consequently, Proposition 5 informs us that a subsequence of u_ε $\boldsymbol{\theta}$-quasi-periodic two-scale converges to some $u \in L^2(\Omega; H^1_{\boldsymbol{\theta}}(Q))$, and moreover $\varepsilon \boldsymbol{\nabla} u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} \boldsymbol{\nabla}_y u$. Let us study the structure of this limit u further. We begin by introducing the densely defined unbounded linear operator $\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}} : H^1_{\boldsymbol{\theta}}(Q) \subset L^2(Q) \to L^2(Q;\mathbb{C}^3)$ which is given by the action

$$w \mapsto \sqrt{a_1}\boldsymbol{\nabla} w, \qquad \text{for } w \in H^1_{\boldsymbol{\theta}}(Q).$$

We now argue that a generalised Weyl's decomposition holds, which was first introduced and proved for the case $\boldsymbol{\theta} = 0$ in [20].

Lemma 3.

$$L^2(Q;\mathbb{C}^3) = \ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big) \oplus \mathrm{Ran}(\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}}).$$

Proof. By the Banach closed ranged theorem, this result will follow if we demonstrate that the range of $\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}}$ is closed, and this fact is implied by Proposition 7.
Indeed, suppose $u_n \in \mathrm{Ran}(\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})$ converges strongly in $L^2(Q;\mathbb{C}^3)$ to some u as $n \to \infty$, i.e. there exists $w_n \in H^1_{\boldsymbol{\theta}}(Q)$ such that $\sqrt{a_1}\boldsymbol{\nabla} w_n$ converges strongly in $L^2(Q;\mathbb{C}^3)$ to u. By Proposition 7, the sequence $w_n^\perp$, where $w_n^\perp$ denotes the orthogonal projection of w_n onto the orthogonal complement $V_{\boldsymbol{\theta}}^\perp$ of $V_{\boldsymbol{\theta}}$ in $H^1_{\boldsymbol{\theta}}(Q)$, is a Cauchy sequence in $H^1_{\boldsymbol{\theta}}(Q)$ and therefore converges, up to some subsequence, to $w \in H^1_{\boldsymbol{\theta}}(Q)$. In particular, $\sqrt{a_1}\boldsymbol{\nabla} w_n = \sqrt{a_1}\boldsymbol{\nabla} w_n^\perp$ converges strongly in $L^2(Q)$ to $\sqrt{a_1}\boldsymbol{\nabla} w$ and, consequently $u = \sqrt{a_1}\boldsymbol{\nabla} w$. Hence, the range of $\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}}$ is closed. □

Let us now describe u in detail.

Lemma 4. *The function u belongs to the Bochner space $L^2(\Omega; V_{\boldsymbol{\theta}})$.*

Proof. Recall that

$$V_{\boldsymbol{\theta}} = \{v \in H^1_{\boldsymbol{\theta}}(Q) \,\big|\, \boldsymbol{\nabla} v = 0 \text{ in } Q_1\} = \ker(\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}}),$$

and so we aim to show that $\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}} u = 0$.
On the one hand we deduce from (6.9) and (5.1) that

$$\lim_{\varepsilon\to 0} \varepsilon \int_\Omega a_1(\tfrac{x}{\varepsilon})\boldsymbol{\nabla} u_\varepsilon \cdot \overline{\phi(x)\boldsymbol{\Phi}(\tfrac{x}{\varepsilon})}\,\mathrm{d}x = 0.$$

Yet, on the other hand, Proposition 3 and the assertion $\varepsilon\boldsymbol{\nabla} u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} \boldsymbol{\nabla}_y u$ imply

$$\begin{aligned}\lim_{\varepsilon\to 0} \varepsilon \int_\Omega a_1(\tfrac{x}{\varepsilon})\boldsymbol{\nabla} u_\varepsilon \cdot \overline{\phi(x)\boldsymbol{\Phi}(\tfrac{x}{\varepsilon})}\,\mathrm{d}x &= \lim_{\varepsilon\to 0} \int_\Omega \varepsilon\boldsymbol{\nabla} u_\varepsilon \cdot \overline{\phi(x) a_1(\tfrac{x}{\varepsilon})\boldsymbol{\Phi}(\tfrac{x}{\varepsilon})}\,\mathrm{d}x \\ &= \int_\Omega \int_Q a_1(y)\boldsymbol{\nabla}_y u(x,y) \cdot \overline{\phi(x)\boldsymbol{\Phi}(y)}\,\mathrm{d}x.\end{aligned}$$

Therefore, as finite sums of $\phi(x)\boldsymbol{\Phi}(y)$ are dense in $L^2(\Omega\times Q;\mathbb{C}^3)$ it follows that $a_1\boldsymbol{\nabla}_y u = 0$ and since $\sqrt{a_1}^{-1} \in L^\infty(Q)$ we find that $u \in L^2(\Omega; V_{\boldsymbol{\theta}})$. □

The following result is of fundamental importance in characterising the ($\boldsymbol{\theta}$-quasi-periodic) limit of the flux $a_1(\frac{\cdot}{\varepsilon})\boldsymbol{\nabla} u_\varepsilon$ in terms of the limit u of the function u_ε. Put another way, this identity is crucial for determining the homogenised coefficients.

Lemma 5. *There exists $\boldsymbol{\xi} \in L^2(\Omega\times Q;\mathbb{C}^3)$ such that, up to a subsequence, $\sqrt{a_1}(\frac{\cdot}{\varepsilon})\boldsymbol{\nabla} u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} \boldsymbol{\xi}$. Moreover, $\boldsymbol{\xi}$ belongs to the Bochner space $L^2\big(\Omega; \ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big)\big)$ and the pair $(u,\boldsymbol{\xi})$ satisfies the identity*

$$\begin{aligned}\int_\Omega\int_Q \boldsymbol{\xi}(x,y)\cdot\overline{\phi(x)\boldsymbol{\Psi}(y)}\,\mathrm{d}y\mathrm{d}x = -\int_\Omega\int_Q \sqrt{a_1}(y)u(x,y)\overline{\boldsymbol{\nabla}_x\phi(x)\cdot\boldsymbol{\Psi}(y)}\,\mathrm{d}y\mathrm{d}x, \\ \forall \phi \in C^\infty(\Omega), \boldsymbol{\Psi} \in \ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big).\end{aligned} \tag{6.10}$$

Proof. By Proposition 2 and (6.9) there exists $\boldsymbol{\xi} \in L^2(\Omega \times Q; \mathbb{C}^3)$ such that, up to a subsequence that we discard, one has

$$\sqrt{a_1}(\tfrac{\cdot}{\varepsilon})\boldsymbol{\nabla} u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} \boldsymbol{\xi}. \tag{6.11}$$

To prove $\boldsymbol{\xi} \in L^2\big(\Omega; \ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big)$, we take in (6.8) test functions of the form $\varepsilon\phi(x)\varphi(\frac{x}{\varepsilon})$, $\phi \in C_0^\infty(\Omega)$, $\varphi \in C^\infty_{\boldsymbol{\theta}}(Q)$, and use (6.9), (6.11) to pass to the limit in ε and deduce that

$$\int_\Omega \int_Q \sqrt{a_1}(y)\boldsymbol{\xi}(x,y) \cdot \overline{\phi(x)\boldsymbol{\nabla}_y \varphi(y)}\, \mathrm{d}y\mathrm{d}x = 0.$$

Therefore, for almost every $x \in \Omega$ one has

$$\int_Q \sqrt{a_1}(y)\boldsymbol{\xi}(x,y) \cdot \overline{\boldsymbol{\nabla}_y \varphi(y)}\, \mathrm{d}y = 0, \qquad \forall \varphi \in H^1_{\boldsymbol{\theta}}(Q),$$

and, hence by Lemma 3 it follows that $\boldsymbol{\xi}(x,y) \in L^2\big(\Omega; \ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big)$.
Let us now prove assertion (6.10). Henceforth, we consider $\Psi \in \ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big)$ to be $\boldsymbol{\theta}$-quasi-periodically extended to $\mathbb{R}^3$. We shall prove below the following "integration by parts" formula:

$$\begin{aligned}\int_\Omega \sqrt{a_1}(\tfrac{x}{\varepsilon})\boldsymbol{\nabla} u_\varepsilon(x) \cdot \overline{\phi(x)\boldsymbol{\Psi}(\tfrac{x}{\varepsilon})}\, \mathrm{d}x = -\int_\Omega \sqrt{a_1}(\tfrac{x}{\varepsilon})u_\varepsilon(x)\overline{\boldsymbol{\nabla}_x \phi(x) \cdot \boldsymbol{\Psi}(\tfrac{x}{\varepsilon})}\, \mathrm{d}x,\\ \forall \phi \in C^\infty(\Omega), \boldsymbol{\Psi} \in \ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big).\end{aligned} \tag{6.12}$$

Using Proposition 3, (6.11) and the convergence $u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} u$, we pass to the limit in the above formula to readily arrive at (6.10).
To prove (6.12), it is sufficient to prove the following: for every $w \in H^1(\mathbb{R}^3)$ one has

$$\int_{\mathbb{R}^3} \sqrt{a_1}(\tfrac{x}{\varepsilon})\boldsymbol{\nabla} w(x) \cdot \overline{\Psi(\tfrac{x}{\varepsilon})}\, \mathrm{d}x = 0. \tag{6.13}$$

Indeed, (6.12) follows from utilising (6.13) and the following facts: for $\phi \in C^\infty(\Omega)$ then $u_\varepsilon\phi$ belongs to $H^1_0(\Omega)$, as $u_\varepsilon \in H^1_0(\Omega)$, and can be trivially extended to $H^1(\mathbb{R}^3)$, and that

$$\int_\Omega \sqrt{a_1}(\tfrac{x}{\varepsilon})\boldsymbol{\nabla} u_\varepsilon(x) \cdot \overline{\phi(x)\Psi(\tfrac{x}{\varepsilon})}\, \mathrm{d}x = \int_\Omega \sqrt{a_1}(\tfrac{x}{\varepsilon})\boldsymbol{\nabla}(u_\varepsilon\overline{\phi})(x) \cdot \overline{\Psi(\tfrac{x}{\varepsilon})}\, \mathrm{d}x - \int_\Omega \sqrt{a_1}(\tfrac{x}{\varepsilon})u_\varepsilon(x)\overline{\boldsymbol{\nabla}_x\phi(x) \cdot \Psi(\tfrac{x}{\varepsilon})}\, \mathrm{d}x.$$

Let us now prove (6.13). For $Q_\varepsilon^{(z)} = \prod_{i=1}^3 \varepsilon(z_i, z_i+1)$, $z \in \mathbb{Z}^3$, it follows that

$$\begin{aligned}\int_{\mathbb{R}^3} \sqrt{a_1}(\tfrac{x}{\varepsilon})\boldsymbol{\nabla} w(x) \cdot \overline{\Psi(\tfrac{x}{\varepsilon})}\, \mathrm{d}x &= \sum_{z\in\mathbb{Z}^3} \int_{Q_\varepsilon^{(z)}} \sqrt{a_1}(\tfrac{x}{\varepsilon})\boldsymbol{\nabla} w(x) \cdot \overline{\Psi(\tfrac{x}{\varepsilon})}\, \mathrm{d}x\\ &= \varepsilon^3 \sum_{z\in\mathbb{Z}^3} \int_Q \sqrt{a_1}(y)\boldsymbol{\nabla} w(\varepsilon y + \varepsilon z) \cdot \overline{\exp(\mathrm{i}\boldsymbol{\theta}\cdot \mathrm{z})\Psi(\mathrm{y})}\, \mathrm{d}y,\end{aligned}$$

where the last equality comes from the change of variables $x = \varepsilon(y+z)$ and recalling that $a_1(y)$ is periodic and Ψ is $\boldsymbol{\theta}$-quasiperiodic. By noting, for $w \in H^1(\mathbb{R}^3)$, that

$$w_\varepsilon(y) := \sum_{z\in\mathbb{Z}^3} w(\varepsilon y + \varepsilon z)\exp(-\mathrm{i}\boldsymbol{\theta}\cdot \mathrm{z}), \qquad \mathrm{y} \in \mathrm{Q},$$

is an element of $H^1_{\boldsymbol{\theta}}(Q)$, and that

$$\boldsymbol{\nabla} w_\varepsilon(y) := \varepsilon \sum_{z\in\mathbb{Z}^3} \boldsymbol{\nabla} w(\varepsilon y + \varepsilon z)\exp(-\mathrm{i}\boldsymbol{\theta}\cdot \mathrm{z}), \qquad \mathrm{y} \in \mathrm{Q},$$

the identity (6.13) follows. □

We are now ready to describe the properties of the macroscopic part of u and express the flux $\boldsymbol{\xi}$ in terms of u.

Lemma 6. *Let $(u, \boldsymbol{\xi})$, $u \in L^2(\Omega; V_{\boldsymbol{\theta}})$ and $\boldsymbol{\xi} \in L^2(\Omega; \ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big)$, be a pair which satisfies the identity (6.10). Then, $u \in U_{\boldsymbol{\theta}}$, see (2.6). That is, for every for $i \in \mathcal{I}_{\boldsymbol{\theta}} = \{i \in \{1,2,3\} \,|\, \theta_i = 0\}$, one has $u = u_i$ on $\Omega \times C_i$, where $\partial_i u_i \in L^2(\Omega)$ with $u_i \nu_i = 0$ on $\partial\Omega$, for $\boldsymbol{\nu}$ the outer unit normal to $\partial\Omega$. Furthermore,*

$$\boldsymbol{\xi}(x,y) = \sqrt{a_1}(y) \sum_{i \in \mathcal{I}_{\boldsymbol{\theta}}} \partial_{x_i} u_i(x) \mathbb{1}_{C_i}(y)[\boldsymbol{\nabla}_y N^{(i)}(y) + \mathbf{e}_i], \qquad x \in \Omega, y \in Q_1. \tag{6.14}$$

Here, $N^{(i)}$ solve (2.8).

The following result immediately follows from the above lemma.

Proposition 8. *For every for $i \in \mathcal{I}_{\boldsymbol{\theta}}$, one has*

$$\int_{C_i} \sqrt{a}_1(y) \boldsymbol{\xi}(x,y)\, \mathrm{d}y = a_i^{\mathrm{hom}} \partial_{x_i} u_i(x) \mathbf{e}_i, \qquad \text{for almost every } x \in \Omega.$$

Here, a^{hom} are given by (2.7), i.e.

$$a_i^{\mathrm{hom}} = \int_{C_i} a_1(y)[\partial_{y_i} N^{(i)}(y) + 1]\, \mathrm{d}y > 0,$$

for $N^{(i)} \in H^1_{\#_i}(C_i) = \{u \in H^1(C_i) \,|\, u \text{ is 1-periodic in the variable } y_i\}$ is the unique solution to the cell problem

$$\begin{cases} \displaystyle\int_{C_i} a_1(y)\big[\boldsymbol{\nabla} N^{(i)}(y) + \mathbf{e}_i\big] \cdot \overline{\boldsymbol{\nabla}\phi}(y)\, \mathrm{d}y = 0, \qquad \forall \phi \in H^1_{\#_i}(C_i), \\ \displaystyle\int_{C_i} N^{(i)} = 0. \end{cases} \tag{6.15}$$

Proof. Equation (6.14) implies

$$\int_{C_i} \sqrt{a}_1(y) \boldsymbol{\xi}(x,y)\, \mathrm{d}y = \partial_{x_i} u_i(x) \int_{C_i} a_1(y)[\boldsymbol{\nabla}_y N^{(i)}(y) + \mathbf{e}_i]\, \mathrm{d}y.$$

For each $j \in \{1,2,3\}\backslash\{i\}$, we set $\phi = y_j$ in (6.15) to determine that

$$\int_{C_i} a_1(y)\big[\boldsymbol{\nabla}_y N^{(i)}(y) + \mathbf{e}_i\big] \cdot \overline{\boldsymbol{\nabla} y_j}(y)\, \mathrm{d}y = \int_{C_i} a_1(y)\big[\boldsymbol{\nabla}_y N^{(i)}(y) + \mathbf{e}_i\big]\, \mathrm{d}y \cdot \mathbf{e}_j = 0.$$

Hence, it follows that

$$\begin{aligned} \int_{C_i} \sqrt{a}_1(y) \boldsymbol{\xi}(x,y)\, \mathrm{d}y &= \partial_{x_i} u_i(x) \int_{C_i} a_1(y)[\boldsymbol{\nabla}_y N^{(i)}(y) \cdot \mathbf{e}_i + 1]\mathbf{e}_i\, \mathrm{d}y \\ &= a_i^{\mathrm{hom}} \partial_{x_i} u_i(x) \mathbf{e}_i, \end{aligned}$$

for almost every $x \in \Omega$. Finally, from (6.15) it follows that

$$\begin{aligned} \int_{C_i} a_1(y)[\boldsymbol{\nabla}_y N^{(i)}(y) \cdot \mathbf{e}_i + 1] &= \int_{C_i} a_1(y)[\boldsymbol{\nabla}_y N^{(i)}(y) + \mathbf{e}_i] \cdot \mathbf{e}_i \\ &= \int_{C_i} a_1(y)[\boldsymbol{\nabla}_y N^{(i)}(y) + \mathbf{e}_i] \cdot \overline{[\boldsymbol{\nabla}_y N^{(i)}(y) + \mathbf{e}_i]}. \end{aligned}$$

Then, the positivity of a_i^{hom} can be seen by the inequality

$$\int_{C_i} a_1(y)[\boldsymbol{\nabla}_y N^{(i)}(y) + \mathbf{e}_i] \cdot \overline{[\boldsymbol{\nabla}_y N^{(i)}(y) + \mathbf{e}_i]} \geq ||a_1^{-1}||^{-1}_{L^\infty(Q_1)} \int_{C_i} |\boldsymbol{\nabla}_y\big(N^{(i)}(y) + y_i\big)|^2\, \mathrm{d}y,$$

and noting that the right-hand side of this inequality can not be zero for this would contract the periodicity of $N^{(i)}$ in the y_i variable. □

Proof of Lemma 6. As $u \in L^2(\Omega; V_{\boldsymbol{\theta}})$, see Lemma 4, then u is constant in each fibre C_i, $i \in \mathcal{I}$. Now if $\theta_i \neq 0$ then u is necessarily zero in C_i. On the other hand, if $\theta_i = 0$, i.e. $i \in \mathcal{I}_{\boldsymbol{\theta}}$, then $u(x,y) = u_i(x)$ for $x \in \Omega$, $y \in C_i$. That is, $u = u_i$ on $\Omega \times C_i$ for some $\mathbf{u} \in L^2(\Omega, L_{\boldsymbol{\theta}})$, where we recall that $L_{\boldsymbol{\theta}}$ is the closed subspace of $\mathbb{C}^3$ spanned by $\{\mathbf{e}_i\}_{i \in \mathcal{I}_{\boldsymbol{\theta}}}$.
Let us now demonstrate that u belongs to the Hilbert space $U_{\boldsymbol{\theta}}$. By substituting $u = u_i$ on $\Omega \times C_i$, $i \in \mathcal{I}$, into (6.10), we deduce that

$$\int_\Omega \int_Q \boldsymbol{\xi}(x,y) \cdot \overline{\phi(x)\Psi(y)}\,\mathrm{d}y\mathrm{d}x = -\sum_{i \in \mathcal{I}_{\boldsymbol{\theta}}} \int_\Omega \int_{C_i} \sqrt{a_1}(y) u_i(x) \overline{\boldsymbol{\nabla}_x \phi(x) \cdot \Psi(y)}\,\mathrm{d}y\mathrm{d}x, \quad \forall \phi \in C^\infty(\Omega), \Psi \in \ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big). \tag{6.16}$$

For fixed $j \in \mathcal{I}_{\boldsymbol{\theta}}$, we will show directly below that there exists a function $\boldsymbol{\Psi}^{(j)} \ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big)$ such that

$$\int_{C_i} \sqrt{a_1}\overline{\boldsymbol{\Psi}^{(j)}} = \mathbf{0} \quad i \neq j, \quad \text{and} \qquad \int_{C_j} \sqrt{a_1}\overline{\boldsymbol{\Psi}^{(j)}} = \mathbf{e}_j. \tag{6.17}$$

Therefore

$$\sum_{i \in \mathcal{I}_{\boldsymbol{\theta}}} u_i(x) \boldsymbol{\nabla}_x \phi \cdot \int_{C_i} \sqrt{a_1}(y) \overline{\boldsymbol{\Psi}^{(j)}(y)}\,\mathrm{d}y = u_j(x) \overline{\partial_{x_j}\phi(x)}, \qquad \text{a.e. } x \in \Omega,$$

and consequently substituting $\boldsymbol{\Psi}^{(j)}$ into (6.16) gives

$$\int_\Omega \left(\int_Q \boldsymbol{\xi}(x,y) \cdot \overline{\boldsymbol{\Psi}^{(j)}}\,\mathrm{d}y \right) \overline{\phi(x)}\mathrm{d}x = -\int_\Omega u_j(x) \overline{\partial_{x_j}\phi(x)}\,\mathrm{d}x, \quad \forall \phi \in C^\infty(\Omega).$$

That is, $\partial_{x_j} u_j(x) = \int_Q \boldsymbol{\xi}(x,y) \cdot \overline{\boldsymbol{\Psi}^{(j)}}\,\mathrm{d}y \in L^2(\Omega)$ and $u_j \nu_j = 0$ on $\partial\Omega$ where $\boldsymbol{\nu}$ is the outer unit normal to $\partial\Omega$, i.e. $u \in U_{\boldsymbol{\theta}}$ if (6.17) holds.
To show (6.17), we note that under the geometric assumptions on cylinders C_i, $i \in \mathcal{I}$, there exists a function

$$\chi_i \in C^\infty(Q) \text{ such that } \chi_i = 1 \text{ on } C_i, \text{ supp}(\chi_i) \text{ compactly contained in } Q \text{ and } \text{supp}(\chi_i) \cap \overline{C_k} = \emptyset \text{ for } k \neq i. \tag{6.18}$$

Then, for each $j \in \mathcal{I}_{\boldsymbol{\theta}} = \{i \in \mathcal{I} \,|\, \theta_i = 0\}$, the function $\boldsymbol{\Psi}^{(j)} = \frac{1}{|C_j|\sqrt{a_1}} \chi_j \mathbf{e}_j$ clearly satisfies (6.17). Furthermore, $\boldsymbol{\Psi}^{(j)}$ belongs to $\ker\big((\sqrt{a_1}\boldsymbol{\nabla})^*\big)$: Indeed, as $\theta_j = 0$, an element $\phi \in H^1_{\boldsymbol{\theta}}(Q)$ is 1-periodic in the variable y_j, and it follows

$$\int_Q \tfrac{1}{\sqrt{a_1}} \chi_j \mathbf{e}_j \cdot \sqrt{a_1}\boldsymbol{\nabla}_y \phi = \int_{C_j} \partial_{y_j}\phi = 0.$$

Therefore, (6.17) holds.
Let us now demonstrate (6.14). For $i \in \mathcal{I}_{\boldsymbol{\theta}}$, and almost every $x \in \Omega$, notice that

$$\sqrt{a_1}(y) u_i(x) \sum_{\substack{j \in \{1,2,3\} \\ j \neq i}} \mathbf{e}_j \partial_{x_j}\phi(x) = \sqrt{a_1}(y) \boldsymbol{\nabla}_y \left(u_i(x) \sum_{\substack{j \in \{1,2,3\} \\ j \neq i}} \partial_{x_j}\phi(x) y_j \right), \quad \text{a.e. } y \in C_i,$$

and, by the geometric assumption of the cylinders, we can extend y_j into Q such that the extensions are elements of $H^1_{\boldsymbol{\theta}}(Q)$ and equal to zero on C_j. Therefore, it follows that

$$\int_\Omega \int_{C_i} \sqrt{a_1}(y) u_i(x) \sum_{\substack{j\in\{1,2,3\}\\ j\neq i}} \partial_{x_j}\phi(x)\Psi_j(y)\,\mathrm{dydx}$$

$$= \int_\Omega \int_Q \sqrt{a_1}(y)\boldsymbol{\nabla}_y \left(u_i(x) \sum_{\substack{j\in\{1,2,3\}\\ j\neq i}} \partial_{x_j}\phi(x) y_j \right) \cdot \boldsymbol{\Psi}(y)\,\mathrm{dydx} = 0.$$

Consequently, (6.16) takes the form

$$\int_\Omega \int_Q \boldsymbol{\xi}(x,y)\cdot \overline{\phi(x)\boldsymbol{\Psi}(y)}\,\mathrm{dyd}x = -\sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}} \int_\Omega \int_{C_i} \sqrt{a_1}(y) u_i(x) \overline{\partial_{x_i}\phi(x)\Psi_i(y)}\,\mathrm{dyd}x,$$
$$\forall \phi \in C^\infty(\Omega), \boldsymbol{\Psi} \in \ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big).$$

Integrating by parts above, which is permissible since $\partial_i u_i \in L^2(\Omega)$, we deduce that

$$\int_\Omega \int_Q \boldsymbol{\xi}(x,y)\cdot \overline{\phi(x)\boldsymbol{\Psi}(y)}\,\mathrm{dyd}x = \sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}} \int_\Omega \int_{C_i} \sqrt{a_1}(y) \partial_{x_i} u_i(x) \overline{\phi(x)\Psi_i(y)}\,\mathrm{dyd}x,$$
$$\forall \phi \in C^\infty_0(\Omega), \boldsymbol{\Psi} \in \ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big).$$

That is, for almost every x, $\boldsymbol{\xi}(x,\cdot)$ is the projection onto $\ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big)$ of the function

$$\mathbf{w}(x,\cdot) = \sqrt{a_1}(\cdot) \sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}} \partial_{x_i} u_i(x) \mathbb{1}_{C_i}(\cdot)\mathbf{e}_i.$$

For each $i \in \mathcal{I}_{\boldsymbol{\theta}}$, let χ_i given by (6.18), and we introduce $\widetilde{N^i} \in H^1(Q)$ the extension into Q, given by Lemma 2, of the function $N^{(i)} \in H^1_{\#_i}(C_i)$ that solves (6.15). It follows that $\sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}} \chi_i \widetilde{N^{(i)}}$ belongs to $H^1_{\boldsymbol{\theta}}(Q)$ and

$$\int_Q a_1 \chi_i [\boldsymbol{\nabla}\widetilde{N^{(i)}} + \mathbf{e}_i]\cdot \overline{\boldsymbol{\nabla}\phi} = \int_{C_i} a_1 [\boldsymbol{\nabla} N^{(i)} + \mathbf{e}_i]\cdot \overline{\boldsymbol{\nabla}\phi} = 0$$

for all $\phi \in H^1_{\boldsymbol{\theta}}(Q)$. That is, $\sqrt{a_1}\chi_i[\boldsymbol{\nabla}\widetilde{N^{(i)}} + \mathbf{e}_i]$ belongs to $\ker\big((\sqrt{a_1}\boldsymbol{\nabla}_{\boldsymbol{\theta}})^*\big)$. Obviously

$$\mathbf{w}(x,y) = \mathbf{w}(x,y) + \sqrt{a_1}(y) \sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}} \partial_{x_i} u_i(x)\chi_i(y)\boldsymbol{\nabla}_y \widetilde{N^{(i)}}(y) - \sqrt{a_1}(y) \sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}} \partial_{x_i} u_i(x)\chi_i(y)\boldsymbol{\nabla}_y \widetilde{N^{(i)}}(y),$$

and

$$\sqrt{a_1}\chi_i \boldsymbol{\nabla}\widetilde{N^{(i)}} = \sqrt{a_1}\boldsymbol{\nabla}(\chi_i \widetilde{N^{(i)}}),$$

since χ_i is piece-wise constant on C. Consequently, as $\boldsymbol{\xi}(x,\cdot)$ is the projection of $w(x,\cdot)$ onto $\ker\big((\sqrt{a_1}\boldsymbol{\nabla})^*\big)$, we have

$$\boldsymbol{\xi}(x,y) = \mathbf{w}(x,y) + \sqrt{a_1}(y) \sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}} \partial_{x_i} u_i(x)\chi_i(y)\boldsymbol{\nabla}_y \widetilde{N^{(i)}}(y) = \sqrt{a_1}(y) \sum_{i\in\mathcal{I}_{\boldsymbol{\theta}}} \partial_{x_i} u_i(x)[\mathbb{1}_{C_i}(y)\mathbf{e}_i + \chi_i(y)\boldsymbol{\nabla}_y \widetilde{N^{(i)}}(y)],$$

Hence, (6.14) holds and the proof is complete.

□

We now conclude with the proof of Theorem 1. That is, we show that u solves (2.9). We being by stating that under the assumption that Ω is star-shaped, standard pull-back and mollification type arguments prove that functions smooth in x are dense in the Hilbert space $U_{\boldsymbol{\theta}}$. Therefore, it is sufficient to show (2.9) holds for such test functions ϕ. Let us take such a ϕ and consider the test functions $\phi_\varepsilon(x) = \phi(x, \frac{x}{\varepsilon})$, $x \in \Omega$ in (6.8). Utilising the convergences

$$u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} u, \qquad \varepsilon \boldsymbol{\nabla} u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} \boldsymbol{\nabla}_y u, \qquad \sqrt{a_1}(\tfrac{\cdot}{\varepsilon}) \boldsymbol{\nabla} u_\varepsilon \overset{2-\boldsymbol{\theta}}{\rightharpoonup} \boldsymbol{\xi},$$

we pass to the limit $\varepsilon \to 0$ in (6.8) to deduce that

$$\int_\Omega \int_C \sqrt{a_1}(y) \boldsymbol{\xi}(x,y) \cdot \overline{\boldsymbol{\nabla}_x \phi}(x,y) \, \mathrm{d}y \mathrm{d}x + \int_\Omega \int_{Q_0} a_0(y) \boldsymbol{\nabla}_y u(x,y) \cdot \overline{\boldsymbol{\nabla}_y \phi}(x,y) \, \mathrm{d}y \mathrm{d}x$$
$$+ \int_\Omega \int_Q u(x,y) \overline{\phi}(x,y) \, \mathrm{d}y \mathrm{d}x = \int_\Omega \int_Q f(x,y) \overline{\phi(x,y)} \, \mathrm{d}y \mathrm{d}x.$$

Then, as $\phi = \phi_i$ on $\Omega \times C_i$, with $\phi \neq 0$ only if $i \in \mathcal{I}_{\boldsymbol{\theta}}$, Proposition 8 implies that

$$\int_\Omega \int_C \sqrt{a_1}(y) \boldsymbol{\xi}(x,y) \cdot \overline{\boldsymbol{\nabla}_x \phi}(x,y) \, \mathrm{d}y \mathrm{d}x = \sum_{i \in \mathcal{I}_{\boldsymbol{\theta}}} \int_\Omega \left(\int_{C_i} \sqrt{a_1}(y) \boldsymbol{\xi}(x,y) \, \mathrm{d}y \right) \cdot \overline{\boldsymbol{\nabla}_x \phi_i}(x) \mathrm{d}x$$
$$= \sum_{i \in \mathcal{I}_{\boldsymbol{\theta}}} \int_\Omega a_i^{\mathrm{hom}} \partial_{x_i} u_i(x) \overline{\partial_{x_i} \phi_i}(x) \mathrm{d}x$$

and (2.9) follows.

Acknowledgements

The author is grateful to Dr Ilia Kamotski and Professor Valery Smyshylaev for many interesting and insightful mathematical discussions connected with two-scale convergence and high-contrast problems. This work was performed under the financial support of the Engineering and Physical Sciences Research Council (Grant EP/M017281/1: "Operator asymptotics, a new approach to length-scale interactions in metamaterials.")

References

[1] G. Allaire, *Homogenization and two-scale convergence*, SIAM J. Math. Anal. **23** (1992), 1482-1518.

[2] A., Avila, G., Griso, B., Miara, *Bandes photoniques interdites en lasticit linarise.* C. R. Math. Acad. Sci. Paris 339, (2004), 377382.

[3] A., Avila, G., Griso, B., Miara, E., Rohan, *Multiscale modeling of elastic waves: theoretical justification and numerical simulation of band gaps.* Multiscale Model. Simul. 7 **1**, (2008), 121.

[4] M. Bellieud, I., Gruais, *Homogenization of an elastic material reinforced by very stiff or heavy fibers. Non-local effects. Memory effects.* J. Math. Pure Appl. **84**, (2005), 5596.

[5] G., Bouchitté, D., Felbacq, *Homogenization near resonances and artificial magnetism from dielectrics.* C. R. Math. Acad. Sci. Paris 339 **5**, (2004), 377382.

[6] M., Briane, *Homogenization of non-uniformly bounded operators: critical barrier for nonlocal effects.* Arch. Ration. Mech. Anal. 164 **1** (2002), 73101.

[7] M., Camar-Eddine, G.W., Milton, *Non-local interactions in the homogenization closure of thermoelectric functionals.* Asymptotic Anal. 41 **34**, (2005), 259276.

[8] M., Camar-Eddine, P., Seppecher, *Determination of the closure of the set of elasticity functionals.* Arch. Ration. Mech. Anal. 170 **3**, (2003), 211245.

[9] M. I., Cherdantsev, *Spectral convergence for high contrast elliptic periodic problems with a defect via homogenization.* Mathematika, Volume 55, Issue 1-2, (2009), pp. 29-57

[10] K. D., Cherednichenko, S., Cooper, S., Guenneau, *Spectral analysis of one-dimensional high-contrast elliptic problems with periodic coefficients.* Multiscale Modeling and Simulation: A SIAM journal, Volume 13, Issue 1, (2015), pp.72-98.

[11] K. D., Cherednichenko, S., Cooper *Homogenisation of the system of high-contrast Maxwell equations.* Mathematika, Volume 61, Issue 2, (2015), pp.475-500.

[12] K. D., Cherednichenko, S., Cooper *Resolvent estimates for high-contrast elliptic problems with periodic coefficients.* Archive for Rational Mechanics and Analysis, Volume 219, Issue 3, (2016), pp.1061-1086.

[13] K. D., Cherednichenko, V. P., Smyshlyaev, V. V., Zhikov, *Non-local homogenised limits for composite media with highly anisotropic periodic fibres.* Proc. R. Soc. Edinb. A 136 **1**, (2006), 87114.

[14] K. D., Cherednichenko, *Two-scale asymptotics for non-local effects in composites with highly anisotropic fibres.* Asymptotic Anal. 49 **1 2**, (2006), 3959.

[15] S., Cooper, *Two-scale homogenisation of partially degenerating PDEs with applications to photonic crystals and elasticity.* PhD Thesis, (2012), University of Bath.

[16] V. N., Fenchenko, E. Ya. Khruslov, *Asymptotic behaviour or the solutions of differential equations with strongly oscillating and degenerating coefficient matrix.* Dokl. Akad. Nauk Ukrain. SSR Ser. A **4** (1980), 2630.

[17] R., Hempel, K., Lienau, *Spectral properties of periodic media in the large coupling limit.* Commun. Part. Diff. Eq. 25 **78**, (2000), 14451470.

[18] V. V. Jikov, S. M. Kozlov, O.A. Oleinik, *Homogenization of Differential Operators and Integral Functionals.*

[19] I. V., Kamotski, V. P., Smyshlyaev, *Localised modes due to defects in high contrast periodic media via homogenization.* Available on ResearchGate: https://www.researchgate.net/publication/253995046_Localised_modes_due_to_defects_in_high_contrast_periodic_media_via_homogenization.

[20] I. V. Kamotski, V. P. Smyshlyaev, *Two-scale homogenization for a class of partially degenerating PDE systems. Preprint arXiv:1309.4579v1,* (2013).

[21] R.V., Kohn, S.P., Shipman, *Magnetism and homogenization of microresonators.* Multiscale Model. Simul. 7 **1**, (2008), 6292.

[22] Z., Liu, C.T., Chan, P., Sheng, *Analytic model of phononic crystals with local resonances.* Phys. Rev. B 71, (2005), 014103.

[23] G, Milton, M., Briane, J., Willis, *On cloaking for elasticity and physical equations with a transformation invariant form.* New J. Phys. 8, (2006), 248.

[24] G., Milton, J., Willis, *On modifications of Newtons second law and linear continuum elastodynamics.* Proc. R. Soc. Lond. A **463**, (2007), 855 880.

[25] G. Nguetseng, *A general convergence result for a functional related to the theory of homogenization*, SIAM J. Math. Anal. **20** (1989), 608-623.

[26] G. V., Sandrakov, *Homogenization of elasticity equations with contrasting coefficients.* Sbornik Math. 190 **12** (1999), 17491806.

[27] V. P., Smyshlyaev, *Propagation and localization of elastic waves in highly anisotropic periodic composites via two-scale homogenization.* Mechanics of Materials, 41, (2009), pp 434-447.

[28] V. V., Zhikov *On an extension of the method of two-scale convergence and its applications*, Sbornik: Mathematics **191**:7 (2000), 973-1014.

[29] V. V., Zhikov *On spectrum gaps of some divergent elliptic operators with periodic coefficients*, St. Petersburg Math. J. **16**:5 (2005), 773-790.

[30] V. V., Zhikov, S. E. Pastukhova *On gaps in the spectrum of the operator of elasticity theory on a high contrast periodic structure.* Journal of Mathematical Sciences, Volume 188, Issue 3, (2013), pp 227-240.